

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/ES 00/00245

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 : C12Q 1/68, C12N 15/62 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 : C12Q, C12N  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practical, search terms used) <b>CIBEPAT, EPODOC, WPI, EMBASE, BIOSIS, MEDLINE, CA, STRAND</b>		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9837235 A (CORNELL RESEARCH FOUNDATION, INC.) 27 August 1998 (27.08.98) page 4, line 35 – page 5, line 15; page 6, line 9- page 7, line 2; page 8, line 3 – page 9, line 3; example 1; claims 1-9	1-9
Y	the whole document	1-9
X	JAOU-CHEM, H et al. «Gene transfer to cultured human endometrial stromal cells: a model to study cyclooxygenase-2 gene regulation», FERTILITY AND STERILITY, 1998, Vol.70, No. 4, pages 734-739. the whole document	1-9
X	HIROYASU, I. et al. «Transcriptional role of nuclear factor KB site in the induction by lipopolysaccharide and suppression by dexamethasone of cyclooxygenase-2 in U937 cell». BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, 1998, Vol. 244, pages 143-148. the whole document	1-9
Y	US 5556754 A (SINGER et al.) 17 September 1996 (17.09.96) the whole document	1-9
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents:</p> <p>“A” document defining the general state of the art which is not considered to be of particular relevance</p> <p>“E” earlier document but published on or after the international filing date</p> <p>“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>“O” document referring to an oral disclosure, use, exhibition or other means</p> <p>“P” document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>“&amp;” document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search 16 October 2000 (16.10.00)		Date of mailing of the international search report 06 November 2000 (06.11.00)
Name and mailing address of the ISA/ <div style="text-align: center;">S.P.T.O.</div>		Authorized officer  Telephone No.



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/ES 00/00245

C. (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5445941 A (YANG) 29 August 1995 (17.09.95) the whole document	1-9
A	US 5569588 A (ASHBY et al.) 29 October 1996 (29.10.96) the whole document	1-9





**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International Application No  
PCT/ES 00/00245

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9837235 A	27.08.1998	AU 6015098 A	09.09.1998
US 5556754 A	17.09.1996	US 5871950 A	16.02.1999
		AU 695120 B	06.08.1998
		WO 9428897 A	22.12.1994
		EP 0702554 A	27.03.1996
		CA 2164641 A	22.12.1994
		AU 056494 A	03.01.1995
		JP 8511266 T	26.11.1996
US 5445941 A	29.08.1995	PL 177706 B	31.01.2000
		IL 109990 A	20.06.1999
		NZ 286125 A	24.11.1997
		AU 2871097 A	25.09.1997
		AU 677319 B	17.04.1997
		ZA 9404160 A	13.12.1995
		PL 303915 A	09.01.1995
		HU 70326 A	28.09.1995
		EP 0629697 A	21.12.1994
		CZ 9401475 A	14.06.1995
		CA 2126294 A	22.12.1994
		BR 9402480 A	25.01.1995
		AU 6470194 A	22.12.1994
		NO 942313 A	22.12.1994
		FI 942958 A	22.12.1994
		CN 1102437 A	10.05.1995
		JP 7184661 A	25.07.1995
US 5569588A	29.10.1996	AU 724474 B	21.09.2000
		JP 10507647 T	28.07.1998
		CA 2202154 A	20.02.1997
		EP 0791078 A	27.08.1997
		AU 6720996 A	05.03.1997
		WO 9706277 A	20.02.1997



(12) SOLICITUD INTERNACIONAL PUBLICADA EN VIRTUD DEL TRATADO DE COOPERACIÓN  
EN MATERIA DE PATENTES (PCT)

(19) Organización Mundial de la Propiedad  
Intelectual  
Oficina internacional



(43) Fecha de publicación internacional  
18 de Enero de 2001 (18.01.2001)

PCT

(10) Número de Publicación Internacional  
**WO 01/04350 A1**

(51) Clasificación Internacional de Patentes: **C12Q 1/68**,  
C12N 15/62

[ES/ES]: Avda. Mare de Deu de Montserrat, 221, E-08041  
Barcelona (ES).

(21) Número de la solicitud internacional: PCT/ES00/00245

(22) Fecha de presentación internacional:  
11 de Julio de 2000 (11.07.2000)

(25) Idioma de presentación: español

(26) Idioma de publicación: español

(30) Datos relativos a la prioridad:  
P 9901557 12 de Julio de 1999 (12.07.1999) ES

(71) Solicitante (para todos los Estados designados salvo  
US): **LABORATORIOS DEL DR. ESTEVE, S.A.**

(72) Inventores: e


(75) Inventores/Solicitantes (para US solamente): **FRESNO  
ESCUDERO, Manuel** [ES/ES]: Centro de Biología  
Molecular "Severo Ochoa", Fac. Ciencias Universidad  
Autónoma de Madrid, Cantoblanco, E-28049 Madrid  
(ES). **IÑIGUEZ PENA, Miguel Angel** [ES/ES]: Centro  
de Biología Molecular "Severo Ochoa", Fac. Ciencias  
Universidad Autónoma de Madrid, Cantoblanco, E-28049  
Madrid (ES).

(74) Mandatario: **CARPINTERO LOPEZ, Francisco**: Her-  
rero & Asociados, S.L., Calle Alcalá, 35, E-28014 Madrid  
(ES).

[Continúa en la página siguiente]

(54) Title: CELL LINE COMPRISING THE PROMOTER OF CYCLOOXYGENASE-2 (COX-2) AND A WITNESS GENE, AND  
USE THEREOF IN THE SEARCH OF COX-2 TRANSCRIPTIONAL INDUCTION SELECTIVE INHIBITORS

(54) Título: LINEA CELULAR QUE COMPRENDE EL PROMOTOR DE LA CICLOOXIGENASA-2 (COX-2) Y UN GEN TES-  
TIGO, Y SU EMPLEO EN LA BUSQUEDA DE INHIBIDORES SELECTIVOS DE LA INDUCCION TRANSCRIPCIONAL DE  
COX-2



(57) Abstract: The cell line comprises a DNA construction which includes all or part of a promoter sequence of the cyclooxygenase-2 (cox-2) gene and a witness gene, operatively jointed to each other, so that said promoter sequence of the cox-2 gene directs the expression of said witness gene in response to an adequate stimulus. The assay method comprises contacting said cell line with the compound to be assayed and determining the existence of a signal which is indicative of the expression of the activity due to the witness gene. Said method is appropriate for searching cox-2 transcriptional induction selective inhibitors through appropriate stimuli.

[Continúa en la página siguiente]



WO 01/04350 A1



(81) **Estados designados (nacional):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) **Estados designados (regional):** patente ARIPO (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), patente euroasiática (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), patente europea (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), patente OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Publicada:**

- Con informe de búsqueda internacional.
- Antes de la expiración del plazo para modificar las reivindicaciones y para ser republicada si se reciben modificaciones.
- Con (una) indicación(es) relativa(s) a material biológico depositado en virtud de lo dispuesto en la Regla 13bis, aparte de la descripción.

Para códigos de dos letras y otras abreviaturas, véase la sección "Guidance Notes on Codes and Abbreviations" que aparece al principio de cada número regular de la Gaceta del PCT.

(57) **Resumen:** La línea celular comprende una construcción de DNA que comprende la totalidad o parte de una secuencia promotora del gen de la ciclooxigenasa 2 (cox-2) y un gen testigo, unidos operativamente entre sí, de manera que dicha secuencia promotora del gen de la cox-2 dirige la expresión de dicho gen testigo en respuesta a un estímulo adecuado. El método de ensayo comprende poner en contacto dicha línea celular con el compuesto a ensayar y determinar la existencia de una señal indicativa de la expresión de actividad debida al gen testigo. Este método es adecuado para la búsqueda de inhibidores selectivos de la inducción a nivel transcripcional de la cox-2 por estímulos apropiados.

LÍNEA CELULAR QUE COMPRENDE EL PROMOTOR DE LA  
CICLOOXIGENASA-2 (COX-2) Y UN GEN TESTIGO, Y SU EMPLEO EN  
LA BÚSQUEDA DE INHIBIDORES SELECTIVOS DE LA INDUCCIÓN  
TRANSCRIPCIONAL DE COX-2

5

**CAMPO DE LA INVENCION**

Esta invención se relaciona, en general, con la  
búsqueda de productos con potenciales aplicaciones  
terapéuticas. En particular, la invención se refiere a un  
10 método para la búsqueda de compuestos inhibidores  
selectivos de la inducción a nivel transcripcional de la  
ciclooxigenasa-2 que comprende el empleo de una línea  
celular que expresa de forma estable una construcción de  
DNA en la que la secuencia promotora del gen de la  
15 ciclooxigenasa-2 dirige la expresión de un gen testigo en  
respuesta a los estímulos apropiados.

**ANTECEDENTES DE LA INVENCION**

La ciclooxigenasa (cox) es una enzima implicada en  
20 numerosos procesos. Se conocen dos isoformas de la cox,  
la ciclooxigenasa 1 (cox-1) y la ciclooxigenasa 2 (cox-  
2). Aunque ambas isoformas están relacionadas con la  
producción de prostaglandinas involucradas en procesos  
fisiológicos, parece ser que la cox-2 es la isoforma  
25 predominante implicada en diversas patologías como la  
inflamación, la carcinogénesis, la angiogénesis y algunos  
procesos neurodegenerativos.

La inducción a nivel transcripcional de la cox-2 se  
produce en respuesta a numerosos factores, entre los que  
30 se encuentran, la expresión de oncogenes, el tratamiento  
con promotores de tumores, mitógenos, estímulos  
proinflamatorios, factores de crecimiento y citoquinas

[revisado en Smith and DeWitt, 1996; Griswold and Adams, 1996; Jouzeau et al., 1997 (véase el apartado relativo a la BIBLIOGRAFÍA)]. En la mayoría de los casos la inducción de esta enzima se traduce en un aumento en la síntesis de prostaglandinas, aunque no se pueden descartar otros modos de actuación.

La capacidad de ciertas drogas de la familia de los antiinflamatorios no esteroideos (NSAIDs) de inhibir la cox-2 explica sus efectos terapéuticos [revisado en Smith and DeWitt, 1996; Griswold and Adams, 1996; Jouzeau et al., 1997]. Asimismo, existen evidencias crecientes de que la inhibición de la cox-2 tanto por NSAIDs como por glucocorticoides o por ciclosporina A posee propiedades inmunosupresoras [Iñiguez et al., 1998; Hall and Wolf, 1997; Zhou et al., 1994; y las revisiones citadas anteriormente]. Otras acciones de la inducción de la cox-2 se refieren a la implicación de ésta en cáncer, angiogénesis y procesos neurodegenerativos como la enfermedad de Alzheimer. Se ha comprobado que tanto la inhibición de la inducción a nivel transcripcional cox-2 como la inhibición enzimática de la cox-2 atenúan estos procesos [Shiff et al., 1996; Tsujii et al., 1997 y 1998; Subbaramiah et al., 1998; Pasinetti, 1998].

Tras el descubrimiento de la isoforma inducible cox-2 de la enzima ciclooxigenasa, los métodos de identificación de nuevos fármacos antiinflamatorios han tenido como objetivo seleccionar compuestos inhibidores selectivos de la actividad enzimática cox-2 frente a la isoforma constitutiva cox-1. Existen varios tipos de sistemas al respecto. Unos utilizan ensayos *in vitro* utilizando enzima cox-2 purificada o semipurificada

[Famaey, 1997; Noreen et al., 1998]. Otros autores utilizan líneas celulares animales o humanas que expresan predominantemente la isoforma cox-2 en condiciones naturales o tras la inducción con estímulos [Famaey, 5 1997; Berg et al., 1997]. Algunos emplean líneas celulares de origen animal o humano en las que sobreexpresan la proteína cox-2 mediante transfección estable del cDNA de la misma [Lora et al., 1997; O'Neill et al., 1995; Cromlish and Kennedy, 1996]. En algún caso 10 se ha llegado a determinar mediante el análisis del mRNA si tales compuestos inhiben a nivel transcripcional la inducción de la cox-2 [Tao et al., 1998; Subbaramiah et al., 1998]. También se han establecido sistemas de estudio de compuestos inhibidores mediante ensayos in 15 vivo, ya sea con sangre entera o con células purificadas de donantes sanos [Famaey, 1997; Brideau et al., 1996].

En cualquier caso, la mayor limitación de estos sistemas radica en que permiten seleccionar compuestos inhibidores de la actividad enzimática de la enzima cox- 20 2, sin atender a sus efectos sobre la inducción de la producción de la proteína, paso previo a la producción de prostaglandinas por esta enzima. Además de esta limitación, ha quedado demostrado que las potencias relativas de estos compuestos varían para el mismo 25 fármaco entre diferentes tipos de ensayos. Asimismo, un aspecto importante a considerar tiene que ver con la inhibición de la actividad cox-2 fisiológica, que también se vería inhibida por el tipo de compuestos identificados por los sistemas mencionados, lo que podría derivar en 30 efectos secundarios adversos.

Existe, por tanto, la necesidad de desarrollar un

método para la búsqueda de compuestos inhibidores selectivos de la inducción a nivel transcripcional de la cox-2 por diferentes estímulos que supere los inconvenientes arriba mencionados.

5

#### COMPENDIO DE LA INVENCION

Esta invención proporciona una solución a la necesidad existente que consiste en el desarrollo de un sistema de ensayo para la búsqueda de compuestos inhibidores selectivos de la inducción a nivel transcripcional de la cox-2 por diferentes estímulos. Este criterio permite seleccionar compuestos que inhiben la producción de la cox-2, con lo que actuarán como inhibidores de las acciones derivadas del incremento de la expresión de la cox-2 y del consiguiente aumento de la producción de prostaglandinas que desencadenan diversos procesos patológicos. Entre otros procesos, se pueden destacar procesos inflamatorios, proliferación celular incontrolada, angiogénesis, carcinogénesis y patologías neurodegenerativas, tal y como se describió anteriormente. El criterio para la selección de compuestos según esta invención radica en la inhibición de la actividad inducible del promotor de la cox-2, con lo cual no se seleccionarán aquellos compuestos que inhiban la actividad basal fisiológica de producción de la cox-2.

Para el desarrollo de la solución aportada por esta invención, ha sido necesario construir una línea celular que expresa de forma estable una construcción de DNA en la que la secuencia promotora del gen de la cox-2 dirige la expresión del gen testigo en respuesta a un estímulo

30



adecuado. La regulación de la expresión del gen de la cox-2 viene determinada por la actividad reguladora de su promotor, mientras que la medida de la actividad del gen testigo proporciona una medida indirecta de la actividad del promotor de la cox-2 en respuesta a diferentes agentes.

Por consiguiente, un objeto de esta invención lo constituye una construcción de DNA (o DNA recombinante) que comprende una secuencia promotora del gen de la cox-2 y un gen testigo, operativamente unidos entre sí, de manera que dicha secuencia promotora del gen dirige la expresión del gen testigo en respuesta a un estímulo adecuado.

Un objeto adicional de esta invención lo constituye un vector, tal como un plásmido o un vector de expresión, que comprende dicha construcción de DNA.

Otro objeto adicional de esta invención lo constituye una línea celular que contiene dicha construcción de DNA, o dicho plásmido que contiene dicha construcción de DNA, y la expresa de forma estable.

Finalmente, otro objeto adicional de esta invención lo constituye un método para la búsqueda de compuestos inhibidores selectivos de la inducción a nivel transcripcional de la ciclooxigenasa-2 que comprende el empleo de dicha línea celular que expresa de forma estable dicha construcción de DNA.

#### BREVE DESCRIPCIÓN DE LAS FIGURAS

La Figura 1 muestra la secuencia de la zona promotora del gen cox-2. Las flechas indican las secuencias de hibridación de los oligonucleótidos

utilizados en la reacción de PCR [reacción en cadena de la polimerasa].

La Figura 2 muestra la estrategia de clonaje de la región promotora del gen *cox-2* en el plásmido pXP2 para obtener la construcción prom2-1906-LUC.

La Figura 3 muestra los resultados del análisis por RT-PCR [transcripción inversa-reacción en cadena de la polimerasa] de la expresión del mRNA de *cox-2* en células Jurkat. En la Figura 3(A) se muestra el efecto del tratamiento con PMA y con PMA + Ionóforo de calcio A23187, en adelante PMA+Ion, en la expresión del mRNA de *cox-2*. En la Figura 3(B) se muestra la inhibición por ciclosporina de la inducción transcripcional de *cox-2*. Como control en ambos casos se muestra el resultado obtenido para los mRNAs no inducibles de la isoforma *cox-1* y de la glicerol-aldehído deshidrogenasa (GAPDH).

La Figura 4 muestra el resultado de la estimulación por PMA o por PMA+Ion de la actividad luciferasa en células Jurkat transfectadas transitoriamente con la construcción prom2-1906-LUC. Como control se comprueba que tanto la construcción prom1-898-LUC como el plásmido vacío pXP2 no son inducibles.

La Figura 5 muestra el resultado de la inhibición por ciclosporina A (CsA) de la estimulación causada por PMA+Ion de la construcción prom2-1906-LUC en la transfección transitoria en células Jurkat.

La Figura 6 muestra los resultados de un experimento de transfección transitoria con la construcción prom2-1906-LUC y tratamiento con dexametasona.

La Figura 7 muestra los resultados de la actividad luciferasa de los diferentes clones obtenidos tras la

transfección estable con la construcción prom2-1906-LUC.

La Figura 8 muestra los resultados de la inhibición por ciclosporina A (CsA) de la estimulación por PMA+Ion de la actividad luciferasa de los clones estables de Jurkat-1906LUC.

La Figura 9 muestra los resultados obtenidos para la inhibición por el glucocorticoide Dexametasona (Dex) sobre la estimulación por PMA+Ion de la actividad luciferasa de los clones estables de Jurkat-1906LUC.

La Figura 10 muestra la inhibición por Resveratrol (Res) de la inducción de la actividad luciferasa de los clones estables como control del sistema de ensayo.

#### DESCRIPCIÓN DETALLADA DE LA INVENCION

Esta invención proporciona una construcción de DNA (o DNA recombinante) que comprende la totalidad o parte de una secuencia promotora del gen de la ciclooxigenasa 2 (cox-2) y un gen testigo, operativamente unidos entre sí, de manera que dicha secuencia promotora del gen de la cox-2 dirige la expresión de dicho gen testigo en respuesta a un estímulo adecuado.

La secuencia promotora del gen cox-2 puede tener cualquier origen, aunque preferentemente dicha secuencia procederá del gen humano de la cox-2.

Como gen testigo puede utilizarse cualquier gen testigo capaz de producir una señal fácilmente detectable, de los habitualmente utilizados en este tipo de ensayos de trasfección, por ejemplo, el gen de la cloranfenicol acetyl transferasa (CAT), el gen de la beta galactosidasa ( $\beta$ -gal), el gen de la luciferasa, por ejemplo, de luciérnaga o de Renilla. En una realización

particular de esta invención dicho gen testigo es el gen de la luciferasa de luciérnaga por la extrema sensibilidad, rapidez, facilidad y bajo coste del ensayo de detección de la misma.

5           La invención también proporciona un vector, tal como un plásmido o un vector de expresión, que contiene la construcción de DNA previamente mencionada. En principio, puede utilizarse cualquier vector apropiado para insertar en él dicha construcción de DNA. Estos vectores son  
10           útiles para transformar células.

          La línea celular proporcionada por esta invención comprende, y expresa de forma estable, dicha construcción de DNA que comprende la totalidad o parte de una secuencia promotora del gen de la cox-2 y un gen testigo,  
15           operativamente unidos entre sí, de manera que dicha secuencia promotora del gen de la cox-2 dirige la expresión de dicho gen testigo en respuesta a un estímulo adecuado.

          La línea celular transformada que contiene la  
20           construcción de DNA previamente mencionada puede proceder de cualquier línea celular adecuada capaz de expresar de forma estable dicha construcción de DNA, por ejemplo, de una línea celular de origen humano tal como una línea de células de tipo linfocito T, células Hep-G2 derivadas de  
25           un carcinoma hepatocelular, células Hela derivadas de un adenocarcinoma de cervix, células tipo monocito-macrófago, por ejemplo, las líneas U937 y THP-1, etc. En una realización particular de esta invención, se ha seleccionado la línea celular Jurkat (descrita  
30           originalmente por Schneider et al., 1977) como ejemplo representativo de una línea celular transformada de tipo

linfocito T como modelo de estudio de la expresión de un gen relacionado con la respuesta inmune. Además, dicha línea celular es fácil de crecer y proporciona un elevado rendimiento de células por unidad de tiempo y por volumen (ml) de cultivo.

La línea celular proporcionada por esta invención puede utilizarse como sistema de ensayo para la búsqueda e identificación (screening) de compuestos inhibidores selectivos de la inducción a nivel transcripcional de la cox-2 por diferentes estímulos.

La línea celular proporcionada por esta invención puede obtenerse fácilmente mediante procedimientos convencionales de Ingeniería Genética, por ejemplo, mediante un procedimiento que comprende (i) el aislamiento de la secuencia promotora del gen de la cox-2, (ii) el clonaje de dicha secuencia en un vector que contiene el gen testigo, en una posición en la que dicha secuencia promotora es capaz de dirigir la expresión de dicho gen testigo, y (iii) la transfección de una línea celular adecuada con dicho plásmido. En el Ejemplo 3 se describe detalladamente una forma concreta de obtener unos clones individuales de células Jurkat transformadas que expresan el gen testigo (luciferasa) de forma estable, denominados Jurkat-C3-1906LUC, Jurkat-F9-1906LUC y Jurkat-C7-1906LUC, en los que se determinó la actividad luciferasa basal y se comprobó que la expresión del gen testigo (luciferasa) se inducía en respuesta a los mismos estímulos que el promotor de la cox-2 tal y como se había establecido previamente con células transfectadas transitoriamente.

El sistema de ensayo (línea celular) proporcionado

por esta invención ha sido validado previamente mediante la transfección transitoria de la construcción prom2-1906-LUC y el análisis de la actividad luciferasa bajo estímulos e inhibidores diferentes. Los resultados obtenidos se compararon con un control no inducible de una construcción similar en la que en lugar del promotor de la cox-2 se sitúa el promotor de la isoforma cox-1 y con el vector vacío pXP2. También se ha procedido a determinar, bajo los mismos estímulos el comportamiento del gen endógeno cox-2 mediante experimentos de RT-PCR en los que se analiza la expresión del mRNA. Como control no inducible se determinó el comportamiento del gen endógeno de la isoforma cox-1 y del gen no inducible de la glicerol-aldehído-fosfato deshidrogenasa (GAPDH). Los principales tratamientos fueron con activadores tales como el éster de forbol PMA (10 ng/ml) y con la combinación de PMA y el Ionóforo de calcio A23187 [PMA+Ion] (1  $\mu$ M). El tratamiento con fármacos inhibidores de la inducción por PMA+Ion se llevó a cabo con dexametasona (1  $\mu$ M), y con ciclosporina A (100 ng/ml).

El Ejemplo 2 incluye unos ensayos de validación del sistema de ensayo proporcionado por esta invención en transfección transitoria, así como de la expresión de los genes endógenos en células Jurkat. También se recogen en los Ejemplos 3 y 4 unos ensayos realizados con los clones estables obtenidos, con compuestos inductores e inhibidores de la inducción del promotor de la cox-2.

El sistema de ensayo proporcionado por esta invención es útil en la búsqueda e identificación (screening) de compuestos inhibidores selectivos de la inducción a nivel transcripcional de la cox-2 por

diferentes estímulos. Este tipo de inhibidores selectivos de la cox-2 puede tener numerosas aplicaciones terapéuticas potenciales ya que las implicaciones derivadas de la inducción de la cox-2 no sólo inciden en la respuesta inflamatoria, sino también en procesos relacionados con la proliferación celular incontrolada y formación de tumores (por ejemplo, la aparición de adenomas, el cáncer de colon y el desarrollo de pólipos y la angiogénesis, entre otros), con acciones inmunosupresoras y con procesos neurodegenerativos como la enfermedad de Alzheimer. Por consiguiente, cabe suponer que los compuestos inhibidores selectivos de la inducción transcripcional de la cox-2 puedan ser útiles como agentes antiinflamatorios, como compuestos capaces de atenuar la proliferación celular incontrolada y/o el proceso de tumorigénesis, como inmunosupresores o como potenciales fármacos con propiedades terapéuticas en la enfermedad de Alzheimer.

La invención también proporciona un método de ensayo para la búsqueda e identificación (screening) de compuestos inhibidores selectivos de la inducción a nivel transcripcional de la cox-2 por un estímulo apropiado (descrito en el Ejemplo 4) que comprende poner en contacto la línea celular proporcionada por esta invención (sistema de ensayo) con el compuesto a ensayar, es decir, con el compuesto cuya potencial actividad inhibidora selectiva de la inducción a nivel transcripcional de la cox-2 se desea ensayar, bajo condiciones que permiten la transcripción de la cox-2, y detectar, y/o medir, la señal indicativa de la expresión de actividad debida al gen testigo. Alternativamente, si

se desea, el método de ensayo objeto de esta invención puede llevarse a cabo poniéndose en contacto la línea celular, el sistema de ensayo y un compuesto activador de la inducción transcripcional de la cox-2.

5 En el método de ensayo proporcionado por esta invención, la regulación de la expresión del gen de la cox-2 viene determinada por la actividad reguladora de su promotor, mientras que la medida de la actividad del gen testigo proporciona una medida indirecta de la actividad  
10 del promotor de la cox-2 en respuesta a diferentes agentes.

El método de ensayo para la búsqueda e identificación de compuestos inhibidores selectivos de la inducción a nivel transcripcional de la cox-2 por un  
15 estímulo apropiado proporcionado por esta invención permite seleccionar compuestos que inhiben la producción de la cox-2 mediante un criterio que radica en la inhibición de la actividad inducible del promotor de la cox-2, con lo cual no se seleccionan aquellos compuestos  
20 que inhiban la actividad basal fisiológica de producción de la cox-2.

Los siguientes ejemplos sirven para ilustrar formas preferidas de realizar la invención sin que deban ser considerados como limitativos del alcance de la misma.

25

#### EJEMPLO 1

Obtención de una construcción de DNA que comprende una  
secuencia del promotor de la cox-2 y el gen de la  
luciferasa

30

##### 1.1 Clonaje del promotor de la cox-2



En primer lugar se procedió al clonaje de la secuencia promotora del gen humano de la cox-2, a partir de la secuencia descrita por Tazawa [Tazawa et al., 1994] representada en la Figura 1.

5 Se utilizó la técnica de la reacción en cadena de la polimerasa (PCR), usando oligonucleótidos cebadores o "primers" diseñados para la amplificación selectiva del fragmento de DNA correspondiente a la secuencia promotora de este gen. Como DNA molde se utilizó DNA genómico de la  
10 línea celular linfocítica humana Jurkat. Los oligonucleótidos utilizados fueron los identificados como SEC.ID.Nº.: 1 y SEC.ID.No.: 2 [véase el apartado relativo a la LISTA DE SECUENCIAS].

Estos oligonucleótidos amplifican una secuencia que  
15 abarca desde el nucleótido -1796 al nucleótido +104 de la zona promotora del gen cox-2 (véase la Figura 1). Para la reacción de PCR se usó el Advantage cDNA PCR kit [Clontech] con 30 ciclos repetitivos de 45 segundos a 94°C y 3 minutos a 68°C en un termociclador PTC-200 [MJ  
20 Research].

### 1.2 Construcción del vector de expresión

Los fragmentos generados tras la amplificación fueron subclonados en el plásmido pXP2 [Nordeen, 1988]  
25 que contiene la secuencia codificante del gen de la luciferasa que se utilizará como gen testigo (véase la Figura 2). Los oligonucleótidos se diseñaron de tal forma que en el extremo 5' contienen una secuencia adicional de reconocimiento por enzimas de restricción. Tras la  
30 amplificación se generan extremos de doble cadena que contienen las dianas de restricción BamHI en el extremo

5' y BglII en el extremo 3'. El vector pXP2 contiene en el sitio múltiple de clonaje una diana BglII, que genera extremos compatibles tanto con extremos BglII como BamHI. Tras la digestión con los enzimas BglII y BamHI del inserto obtenido por PCR conteniendo la secuencia promotora, y del vector pXP2 con BglII, se procedió a la ligación de estas secuencias. De este modo se obtuvo el plásmido prom2-1906-LUC en el que la secuencia (-)1796-(+)104 del promotor de la cox-2 se sitúa por delante del gen de la luciferasa, dirigiendo su expresión. Esta construcción se secuenció para comprobar la fidelidad de la secuencia del promotor y verificar el sitio de clonaje.

15

### Ejemplo 2

20

Experimentos de análisis de la regulación de la actividad del promotor de la cox-2 mediante experimentos de RT-PCR y de transfección transitoria de la construcción prom2-1906-LUC en la línea celular Jurkat

25

30

Con el fin de estudiar la regulación de la expresión del gen cox-2 endógeno en la línea celular Jurkat se procedió al análisis de la expresión de su mRNA mediante experimentos de RT-PCR. A su vez, con el fin de validar la construcción prom2-1906-LUC y comprobar que la regulación del promotor clonado se corresponde con la esperada, se realizaron experimentos de transfección transitoria de la construcción prom2-1906-LUC utilizando la línea celular Jurkat. En ambos experimentos las células se trataron con compuestos estimuladores e inhibidores de la inducción del promotor de cox-2.

El tratamiento con compuestos activadores se llevó a cabo utilizando el éster de forbol PMA (Phorbol 12-Myristate 13-Acetate) (10 ng/ml) (Sigma) y la combinación de PMA y el ionóforo de Calcio A23187 (1 $\mu$ M) (Sigma), en  
5 adelante PMA+Ion.

El tratamiento con fármacos inhibidores de la inducción por PMA+Ion se llevó a cabo con ciclosporina A (CsA) (100 ng/ml), o el glucocorticoide sintético Dexametasona (Dex) (1 $\mu$ M) (Sigma).

10

#### 2.1 Regulación de la expresión del mRNA de cox-2 en células Jurkat

Los resultados obtenidos del análisis por RT-PCR de la expresión del mRNA de cox-2 en células Jurkat se muestran en la Figura 3, donde puede observarse lo  
15 siguiente:

a) el tratamiento con PMA (10 ng/ml) produce un ligero incremento de la expresión del mRNA de cox-2, mientras que el tratamiento combinado con PMA+Ion produce  
20 el mayor aumento en la expresión de este gen a nivel transcripcional [Figura 3(A)]; y

b) la inhibición por CsA (100 ng/ml) de la inducción transcripcional de cox-2 [Figura 3(B)].

Como control en ambos casos se muestra el resultado  
25 obtenido para los mRNAs no inducibles de la isoforma cox-1 y de la glicerol aldehído fosfato deshidrogenasa (GAPDH).

#### 2.2 Evaluación de la actividad luciferasa bajo diferentes estímulos

  
30

Se ha analizado la actividad luciferasa en células

Jurkat transfectadas transitoriamente con la construcción prom2-1906-LUC utilizando el agente Lipofectina [Life Technologies]. Las células fueron estimuladas con PMA (10 ng/m) (Sigma) y con la combinación PMA+Ion (1  $\mu$ M) (Sigma).

Como control no inducible se utilizó una construcción similar en la que en lugar del promotor de la cox-2 se situó el promotor de la isoforma cox-1 [prom1-898-LUC] y con el vector vacío pXP2.

La actividad luciferasa se determinó utilizando el kit "Luciferase Assay System" [Promega], con  $1 \times 10^6$  células que se lisaron en 50  $\mu$ l de tampón de lisis. En los extractos obtenidos se midió la emisión de luz producida con un luminómetro Monolight 2010 [Analytical Luminiscence Laboratory] con un sistema de inyección automática de 100  $\mu$ l de reactivo.

Los resultados obtenidos se muestran en la Figura 4. Como puede apreciarse en dicha figura, los resultados obtenidos en este ensayo son comparables a los obtenidos en el análisis del mRNA de cox-2 [Ejemplo 2.1], es decir, el tratamiento con PMA+Ion produce el mayor aumento en el número de veces de inducción de la actividad luciferasa (aproximadamente 12 veces la basal). Estos datos demuestran que la secuencia promotora clonada se comporta de forma similar al gen endógeno. Como control se comprueba que tanto la construcción prom1-898-LUC como el plásmido vacío pXP2 no son inducibles.

### 2.3 Evaluación de la actividad luciferasa bajo diferentes inhibidores

Se ha analizado la inhibición de la estimulación por

PMA (10 ng/ml) (Sigma) o PMA+Ion (1  $\mu$ M) (Sigma) sobre la actividad luciferasa en células Jurkat transfectadas transitoriamente con la construcción prom2-1906-LUC, mediante el empleo del fármaco inmunosupresor ciclosporina A (CsA) (100 ng/ml). Los resultados obtenidos se muestran en la Figura 5 donde se pone de manifiesto que el tratamiento con CsA (100 ng/ml) disminuye la estimulación del promotor de la cox-2 en respuesta a PMA+Ion hasta valores similares a los basales.

Asimismo, se ha analizado la inhibición de la estimulación por PMA+Ion (1  $\mu$ M) (Sigma) sobre la actividad luciferasa en células Jurkat transfectadas transitoriamente con la construcción prom2-1906 mediante el glucocorticoide dexametasona (1  $\mu$ M) (Sigma). Los resultados obtenidos se muestran en la Figura 6 donde se pone de manifiesto que el tratamiento con dexametasona (1  $\mu$ M) (Dex) disminuye la estimulación del promotor de cox-2 en respuesta a PMA+Ion.

### Ejemplo 3

Obtención de una línea celular que expresa de forma estable una construcción de DNA que comprende una secuencia promotora de cox-2 y el gen de la luciferasa.

Para la creación de la línea celular que expresa de forma estable la construcción prom2-1906-LUC, se cotransfectaron células Jurkat con el vector prom2-1906-LUC y un vector denominado pcDNA3.1/Hygro (Invitrogen) que contiene el gen de resistencia a la higromicina. La transfección se realizó mediante la técnica de

electroporación en cubetas de 0,4 cm (BioRad) con  $15 \times 10^6$  células en 0,5 ml de medio completo [medio RPMI suplementado con 10% de suero fetal, L-Glutamina 2 mM y una mezcla de antibióticos] (Todos estos productos fueron adquiridos a Life Technologies). Las células se incubaron en hielo durante 10 minutos con 25  $\mu$ g del plásmido prom2-1906-LUC y 5  $\mu$ g del vector pCDNA3.1./Hygro. Tras este periodo, las células se electroporaron en un aparato Gene Pulser II (BioRad) a 1.500  $\mu$ Faradios de capacitancia y 280 Voltios de corriente. A continuación, las células se incubaron en hielo durante 10 minutos antes de añadir 10 ml de medio completo. Las células se cultivaron en este medio en frascos de cultivo (Nunc) de 75 cm<sup>2</sup> durante 48 horas en un incubador de células a 37°C, con una humedad del 95% y 5% de CO<sub>2</sub>. En este momento se cambió el medio por medio completo sin antibióticos al que se añadió higromicina (Boehringer Mannheim) a una concentración de 200  $\mu$ g/ml. Las células se cultivaron en este medio durante 30 días con sucesivos cambios de medio. Durante este periodo se selecciona la población resistente que sobrevive al tratamiento con el antibiótico selectivo, es decir los transfectantes estables para el gen de resistencia al antibiótico higromicina. En esta población se analizó la expresión del gen de la luciferasa con el fin de determinar la presencia de transfectantes estables para el plásmido prom2-1906-LUC. Para ello,  $1 \times 10^6$  células se lisaron en 50  $\mu$ l de tampón de lisis (Promega) y con los extractos obtenidos se procedió a determinar la actividad luciferasa con los reactivos contenidos en el kit de "Luciferase Assay System" [Promega]. La medición de la emisión de luz producida se determinó en un

luminómetro MonoLight 2010 (Analytical Luminiscence Laboratory) con un sistema de inyección automática de 100  $\mu$ l de reactivo.

De esta población policlonal (Jurkat-pool-1906LUC) se realizó una dilución límite en placas de 96 pocillos en medio completo con higromicina con el fin de obtener clones individuales que expresaran el gen de la luciferasa de forma estable. Estos clones se crecieron hasta obtener al menos  $1 \times 10^6$  células con las que realizar la medición de la actividad luciferasa tal como se ha descrito anteriormente. De esta forma se obtuvieron tres clones individuales denominados Jurkat-C3-1906LUC, Jurkat-F9-1906LUC y Jurkat-C7-1906LUC. En estos clones se determinó la actividad luciferasa basal en RLUs (unidades relativas de luminiscencia) y se comprobó que la expresión del testigo luciferasa se inducía en respuesta a los mismos estímulos que el promotor de cox-2 tal y como se había establecido previamente con células transfectadas de forma transitoria (véase la Figura 7). En los tres clones, los valores basales de actividad luciferasa se incrementan de 3 a 6 veces con un tratamiento de 6 horas con el éster de forbol PMA y hasta 10 - 20 veces con un tratamiento combinado de PMA+Ion, de forma similar a los resultados obtenidos en las transfecciones transitorias.

#### Ejemplo 4

Establecimiento de un sistema de ensayo de compuestos que regulan la expresión del gen cox-2 en los clones de la línea celular que expresa de forma estable una construcción de DNA que comprende una secuencia promotora de cox-2 y el gen de la luciferasa.

Los clones se cultivaron en placas de 96 pocillos a una densidad de  $1 \times 10^5$  células en 200  $\mu$ l de medio RPMI suplementado con 2% de suero fetal, L-Glutamina 2 mM y una mezcla de antibióticos. Las células se trataron durante 6 horas con diferentes concentraciones de los compuestos cuya actividad se pretende ensayar. En el caso del ensayo de la actividad de estos compuestos sobre la inducción de la actividad del promotor de cox-2, las células se trataron con PMA+Ion durante 5 horas, tras 1 hora de pretratamiento con el compuesto a ensayar. Tras este periodo las células se lisaron en 50  $\mu$ l de tampón de lisis y se determinó su actividad luciferasa utilizando 20  $\mu$ l en un luminómetro tal y como se describió en el Ejemplo 2.2. A continuación se muestran algunos resultados obtenidos con compuestos previamente descritos como inhibidores de la estimulación del promotor de cox-2.

La Figura 8 muestra los resultados obtenidos con el compuesto ciclosporina A (CsA) el cual produce una inhibición de la estimulación obtenida con PMA+Ion en los clones estables de Jurkat, de forma similar a lo previamente descrito (Iñiguez et al., 1998), y a lo observado en las transfecciones transitorias ilustradas en el Ejemplo 2.3.

La Figura 9 muestra los resultados obtenidos con el compuesto Dexametasona (Dex), que, como glucocorticoide y anti-inflamatorio, produce una inhibición de la estimulación obtenida con PMA+Ion en los clones estables de Jurkat, como corresponde a lo ya descrito (Smith and DeWitt, 1996) y a lo observado previamente en las transfecciones transitorias del Ejemplo 2.3.



La Figura 10 muestra los resultados obtenidos con el compuesto Resveratrol (Res), descrito recientemente como inhibidor de la estimulación por PMA del gen cox-2 (Subbaramiah, et al., 1998). Este compuesto produce una inhibición de la estimulación obtenida con PMA+Ion en los clones estables de Jurkat.

De esta forma queda demostrada la validez del sistema de ensayo por el comportamiento similar de los clones obtenidos, de las transfecciones transitorias y de los resultados obtenidos con el mRNA endógeno. Con el uso de compuestos cuya actividad sobre el promotor de cox-2 es conocida, ya sean estimuladores o inhibidores, queda establecido que es posible detectar compuestos no descritos previamente que regulen tanto positivamente como negativamente la expresión basal o inducida del gen cox-2 con el sistema de ensayo desarrollado en la presente invención.

#### DEPOSITO DEL MATERIAL BIOLÓGICO

Una muestra de una línea celular Jurkat, denominada J-1906-F9, que expresa de forma estable una construcción de DNA que comprende una secuencia promotora del gen de la cox-2 y el gen de la luciferasa, ha sido depositada en la European Collection of Animal Cell Cultures (ECACC) [Salisbury, R. Unido] el 24 de marzo de 1999 y ha recibido el número de acceso ECACC 99032405.

Una muestra del plásmido prom2-1906-LUC, insertado en *Escherichia coli* DH5, denominada DH5 prom2-1906-LUC, ha sido depositada en la Colección Española de Cultivos Tipo (CECT) [Burjassot, Valencia] el 24 de marzo de 1999 y ha recibido el número de acceso CECT 5145.

## BIBLIOGRAFÍA

Berg, J., Christoph, T., Widerna, M., and Bodenteich, A. 1997. Isoenzyme-specific cyclooxygenase inhibitors: a whole cell assay system using the human erythroleukemic cell line HEL and the human monocytic cell line Mono Mac 6. *J. Pharmacol. Toxic Methods*. 37: 179-86.

Brideau, C., Kargman, S., Liu S., Dallob A.L., Enrich E.W., Rodger, I.W., And Chan C.C. 1996. A human whole cell assay for clinical evaluation of biochemical efficacy of cyclooxygenase inhibitors. *Inflamm. Res*. 45: 68-74.

Cromlish W.A., and Kennedy, B.P. 1996. Selective inhibition of cyclooxygenase-1 and -2 using intact insect cell assays. *Biochem. Pharmacol*. 52: 1777-85.

Famaey, J.P. 1997. In vitro and in vivo pharmacological evidence of selective cyclooxygenase-2 inhibition by nimesulide: an overview. *Inflamm. Res*. 46: 437-446.

Griswold, D.E., and Adams, J.L. 1996. Constitutive cyclooxygenase (cox-1) and inducible cyclooxygenase (cox-2): Rationale for selective inhibition and progress to date. *Med. Res. Rev*. 16: 181.

Hall, V.C., and Wolf, R.E. 1997. Effect of Tenidap and nonsteroidal antiinflammatory drugs on the response of cultured human T cells to interleukin 2 in rheumatoid arthritis. *J. Rheumatol*. 24:1467.

Iñiguez, M.A., Punzón, C., and Fresno, M. 1998. Induction of cyclooxygenase-2 on activated T lymphocytes; regulation of T cell activation by cyclooxygenase-2 inhibitors. Enviado a publicar.

Jouzeau, J.Y., Terlain, B., Abid, A., Nedelec, E., and Netter, P. 1997. Cyclooxygenase isoenzymes. How recent findings affect thinking about nonsteroidal anti-inflammatory drugs. *Drugs* 53: 563.

5 Lora, M., Morisset, S., Menard, H.A., Leduc, R., and de Brum-Fernandes, A.J. 1997. Expression of recombinant human cyclooxygenase isoenzymes in transfected COS-7 cells in vitro and inhibition by tenoxicam, indomethacin and aspirin. *Prostaglandins Leukot. Essent. Fatty Acids*.  
10 56: 361-7.

Nordeen, S.K. 1988. Luciferase reporter gene vectors for analysis of promoters and enhancers. *Biotechniques* 6: 454-457.

15 Noreen, Y., Ringbom, T., Perera, P., Danielson, H., And Bohlin, L. 1998. Development of a radiochemical cyclooxygenase-1 and -2 in vitro assay for identification of natural products as inhibitors of prostaglandin biosynthesis. *J. Nat. Prod.* 61: 2-7.

20 O'Neill, G.P., Kennedy, B.P. Mancini, J.A., Kargman, S., Ouellet, M., Yergey, J., Falgoutret, J.P., Cromlish, W.A., Payett, P., Chan, C.C. et al. 1995. Selective inhibitors of Cox-2. *Agents Actions Suppl.* 46: 159-68.

25 Pasinetti, G.M. 1998. Cyclooxygenase and inflammation in Alzheimer's disease. Experimental approaches and clinical interventions. *J. Neurosci. Res.* 54: 1-6.

30 Schneider U., Schwenk, H.U., and Bornkamm, G. 1977. Characterization of EBV-genome negative "null" and "T" cell lines derived from children with acute lymphoblastic leukemia and leukemic transformed non-Hodkin lymphoma. *Int. J. Cancer*, 19:521-6.

Sheng, H., Shao, J., Kirkland, S.C., Isakson, P., Coffey, R.J., Morrow, J., Beauchamp, R.D., and DuBois, R.N. 1997. Inhibition of human colon cancer cell growth by selective inhibition of cyclooxygenase-2. *J. Clin. Invest.* 99: 2254-2259.

Shiff, S.J., Koutsos, M.I., Qiao, L., and Rigas, B. 1996. Nonsteroidal antiinflammatory drugs inhibit the proliferation of colon adenocarcinoma cells: effect on cell cycle and apoptosis. *Exp. Cell Res.* 222, 179-188.

Smith, W.L., and DeWitt, D.L. 1996. Prostaglandin Endoperoxide H Synthase-1 and -2. *Adv. Immunol.* 62:167.

Subbaramiah, K., Chung, W.J., Michualart, P., Telang, N., Tanabe, T., Inoue, H., Jang, M., Pezzuto, J.M., and Dannenberg, A.J. 1998. Resveratrol inhibits cyclooxygenase-2 transcription and activity in phorbol-ester treated human mammary epithelial cells. *J. Biol. Chem.* 273: 21875-882.

Tao, X., Schulze-Koops, H., Ma, L., Cai, J., Mao, Y., and Lipsky, P.E. 1998. Effects of *Tripterygium wilfordii* hook extracts on induction of cyclooxygenase 2 activity and prostaglandin E2 production. *Arthritis Rheum.* 41: 130-8.

Tazawa, R., Xu, X.M., Wu, K.K., and Wang, L.H. 1994. Characterization of the genomic structure, chromosomal location and promoter of human prostaglandin H synthase-2 gene. *Biochem. Biophys. Res. Comm.* 203:190-194.

Tsujii M., Kawano, S., Tsuji, S., Sawaoka, H., Matsatsugu, H., and DuBois, R.N. 1998. Cyclooxygenase regulates angiogenesis induced by colon cancer cells. *Cell* 93: 705-716.

Tsujii, M., Kawano, S., and DuBois, R.N. 1997.

Cyclooxygenase-2 expression in human colon cancer cells increases metastatic potential. *Proc. Natl. Acad. Sci. USA* 94: 3336-40.

5        Zhou, L., Ritchie, D., Wang, E.Y., Barbone, A.G.,  
Argentier D., And Lau, C.Y. 1994. Tepoxalin, a novel  
immunosuppressive agent with a different mechanism of  
action from cyclosporin A. *J. Immunol.* 153:5026.

## REIVINDICACIONES

1. Una construcción de DNA que comprende la totalidad o parte de una secuencia promotora del gen de la ciclooxigenasa 2 (cox-2) y un gen testigo, operativamente unidos entre sí, de manera que dicha secuencia promotora del gen de la cox-2 dirige la expresión de dicho gen testigo en respuesta a un estímulo adecuado.

10

2. Construcción según la reivindicación 1, en la que dicha secuencia promotora del gen cox-2 procede del gen humano de la cox-2.

15

3. Construcción según la reivindicación 2, en la que dicha secuencia promotora del gen cox-2 está constituida por la secuencia comprendida entre el nucleótido (-)1796 y el nucleótido (+)104 del promotor de la cox-2 humana.

20

4. Construcción según la reivindicación 1, en la que dicho gen testigo se selecciona entre el gen de la luciferasa, el gen de la cloranfenicol acetil transferasa y el gen de la beta galactosidasa.

25

5. Un vector que comprende una construcción de DNA según cualquiera de las reivindicaciones 1 a 4.

30

6. Una línea celular que comprende una construcción según cualquiera de las reivindicaciones 1 a 4 o transformada con un vector según la reivindicación 5.

7. Una línea celular según la reivindicación 6, en la que dicha línea celular deriva de una línea celular de origen humano.

5           8. Una línea celular según la reivindicación 7, en la que dicha línea celular de origen humano es una línea de células Jurkat.

10           9. Método de ensayo para la búsqueda de compuestos inhibidores selectivos de la inducción a nivel transcripcional de la ciclooxigenasa-2 por un estímulo apropiado, que comprende poner en contacto una línea celular según cualquiera de las reivindicaciones 6 a 8, con un compuesto cuya potencial actividad inhibidora  
15           selectiva de la inducción a nivel transcripcional de la cox-2 se desea ensayar, bajo condiciones que permiten la transcripción de la cox-2, y detectar, y/o medir, la señal indicativa de la expresión de actividad debida al gen testigo.





1/10

-1841 GAATTCAGGATTGTAATGTAATAATTTAGTACTCTCTCACAGTATGGATTCTAACATGGCTTCTAACCCAACTAACATTAGTAGCTCTAATATAAACT  
Oligonucleotide n° 1  
-1741 TCAATTTTCAGTAGATGCAACCTACTCCTTTAAATGAAACAGAAGATTGAAATTATTAATATCAAAAAGAAAATGATCCACGCTCTTAGTTGAAATT  
-1641 TCATGTAAGATTCCATGCAATAAATAGGAGTCCATAAATGGAATGATGAAATATGACTAGAGGAGGAGAAAGGCTTCCTAGATGAGATGGAATTTTAGT  
-1541 CATCCGTGTCTCATGAAGAATCAGATGTGTACACTAAGCAAAACAGTTAAAAAAAACCTCCAAGTGAGTCTTATTTATTTTTTCTFATAAGACTT  
-1441 CTACAAATTGAGGTACCTGGTGTAGTTTATTTTCAGGTTTATGCTGTCAATTTCTGTAAATGCTAAGGACTTAGGACATACTGAATTTCTATTTTCC  
-1341 ACTTCTTTTCTGGTGTGTGTATATATATATGTATATATACACACACACATACATATATATATTTTAGTATCTCACCTCACATGCTCCTCCCTGA  
-1241 GCACTACCCATGATAGATGTTAAACAAAAGCAAAGATGAAATTCCAACTGTAAATCTCCCTTCCATCTAATTAATTCCTCATCCAATATGTCCAAA  
-1141 ACGAGAATAGAAAATTAGCCCCAATAAGCCCAGGCAACTGAAAAGTAAATGCTATGTTGACTTTGATCCATGGTCACAACCTCATAATCTGGAAAAGTG  
-1041 GACAGAAAAGACAAAAGAGTGAACTTTAAACTCGAATTTATTTTACCAGTATCTCCTATGAAGGGCTAGTAACCAAAATAATCCACGCATCAGGGAGAG  
-941 AAATGCCTTAAGGCATACGTTTGGACATTTAGCGTCCCTGCAATTTCTGGCCATCCCGCTTCCTTTGTCATCAGAAGGCAGGAACTTTATATTGGT  
- 841 GACCCGTGGAGCTCACATTAATACTATTACAGGGTAACTGCTTAGGACCAGTATTATGAGGAGGATTACCTTTCCCGCTCTCTTCCAAGAAACAAGGA  
-741 GGGGGTGAAGGTACGGAGAACAGTATTTCTTCTGTTGAAAGCAACTTAGCTACAAAGATAAATTACAGCTATGTACACTGAAGGTAGCTATTTTCATTCCA  
-641 CAAAATAAGAGTTTTTTTAAAAAGCTATGTATGTATGTCTGCATATAGAGCAGATATACAGCCTATTAAGCGTCGTCACTAAACATAAAACATGTCAGC  
-541 CTTTCTTAACCTTACTCGCCCCAGTCTGTCCCGACGTGACTTCCTCGACCCTCTAAAGACGTACAGACCAGACACGGCGCGCGGGGAGAGGGGATT  
-441 CCCTGCGCCCCCGGACCTCAGGGCCGCTCAGATTCTTGAGAGGAAGCCAAGTGCTTCTGCCCCCCCCGGTATCCCATCCAAGGCGATCAGTCCAGA  
-341 ACTGGCTCTCGGAAGCGCTCGGGCAAAGACTGCGAAGAAGAAAAGACATCTGCGGAAACCTGTGCGCTGGGGCGGTGGAACCTGGGGAGGAGAGGGAG  
-241 GGATCAGACAGGAGAGTGGGGACTACCCCTCTGCTCCCAAATTGGGGCAGCTTCTGGGTTTCCGATTTTCTCATTTCCGTGGGTAAAAACCTGCC  
-141 CCACGGGGCTTACGCAATTTTTTAAGGGGAGAGGAGGAAAAATTTGIGGGGGTACGAAAAGCGGAAAAGAACAGTCATTTCTGCATGCGCTTGG  
-41 TTTTCAGTCTTATAAAAAGGAAGGTCTCTCGGTAGCGACCAATTGTCATACGACTGCACTGAGCGTCAGGAGCACGTCCAGGAACCTCAGCAGCG  
+60 CCTCCTTCAGCTCCACAGCCAGACGCCCTCAGACAGCAAGCCTACCCCCGCGCGCGCCCTGCCCGCCGCTGCGATG  
Oligonucleotide n° 2

FIG. 1



2/10




FIG. 2



3/10




FIG. 3



4/10




FIG. 4





5/10




FIG. 5



6/10




FIG. 6



7/10




FIG. 7



8/10




FIG. 8





9/10




FIG. 9



10/10




FIG. 10



## LISTA DE SECUENCIAS

## (1) INFORMACION GENERAL:

## (i) SOLICITANTE:

- 5 (A) NOMBRE: Laboratorios del Dr. Esteve, S.A.  
(B) DIRECCION: Avda. Mare de Deu de Montserrat,  
221  
(C) CIUDAD: Barcelona  
(E) PAIS: España  
10 (F) CODIGO POSTAL: 08041  
(G) TELEFONO: 93 446 60 00  
(H) TELEFAX: 93 450 32 02

## (ii) TITULO DE LA INVENCION:

- 15 LINEA CELULAR QUE COMPRENDE EL PROMOTOR DE LA  
CICLOOXIGENASA-2 (COX-2) Y UN GEN TESTIGO, Y SU  
EMPLEO EN LA BÚSQUEDA DE INHIBIDORES SELECTIVOS DE  
LA INDUCCIÓN TRANSCRIPCIONAL DE COX-2  
20

## (iii) NUMERO DE SECUENCIAS: 2

## (iv) FORMA LEGIBLE POR ORDENADOR:

- 25 (A) MEDIO: Diskete  
(B) ORDENADOR: IBM PC compatible  
(C) SISTEMA OPERATIVO: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version  
#1.30 (EPO)

- 30 (2) INFORMACION DE LA SECUENCIA IDENTIFICADA N°  
[SEC.ID.N°]: 1:



## (i) CARACTERISTICAS DE LA SECUENCIA:

- (A) LONGITUD: 33 nucleótidos
- (B) TIPO: ácido nucleico
- (C) N° DE CADENAS: monocatenaria
- (D) TOPOLOGIA: lineal

## (ii) TIPO DE MOLECULA: DNA

## (xi) DESCRIPCION DE LA SECUENCIA: SEC. ID. N°: 1:

GGGGGATCCG GATTCTAACA TGGCTTCTAA CCC 33

## 10 (2) INFORMACION PARA LA SECUENCIA IDENTIFICADA N°: 2:

## (i) CARACTERISTICAS DE LA SECUENCIA:

- (A) LONGITUD: 30 nucleótidos
- (B) TIPO: ácido nucleico
- (C) N° DE CADENAS: monocatenaria
- (D) TOPOLOGIA: lineal

## (ii) TIPO DE MOLECULA: DNA

## (xi) DESCRIPCION DE LA SECUENCIA: SEC. ID. N°: 2:

GGGAGATCTG GTAGGCTTTG CTGTCTGAGG 30





PCT

2000186

Original (para PRESENTACION) - impreso el 05.07.2000 02:10:05 PM

0-1	Formulario - PCT/RO/134 (EASY) Indicaciones relativas al Depósito de Microorganismo(s) u otro Material Biológico (Regla 13bis del PCT)	
0-1-1	Preparado usando	PCT-EASY Version 2.90 (actualizado el 10.05.2000)
0-2	Solicitud internacional No..	<b>PCT/ E S 00 / 00 2 4 5</b>
0-3	Referencia al expediente del solicitante o del mandatario	2000186
1	Las indicaciones hechas a continuación se refieren al(los) microorganismo(s) u otro material biológico depositado mencionado en la descripción en:	
1-1	página	21
1-2	línea	11-24
1-3	Identificación del depósito	
1-3-1	Nombre de la institución de depósito	European Collection of Cell Cultures Vaccine Research and Production Laboratory, Public Health Laboratory Service, Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom
1-3-2	Dirección de la institución depositaria	24 Marzo 1999 (24.03.1999)
1-3-3	Fecha de depósito	ECACC 99032405
1-3-4	Número de orden	
1-4	Indicaciones adicionales	Depósito de Línea Celular JURKAT
1-5	Las indicaciones son dadas para los siguientes Estados designados	todos los Estados designados
1-6	Suministro separado de indicaciones Estas indicaciones serán presentadas a la Oficina Internacional posteriormente	NINGUNA
2	Las indicaciones hechas a continuación se refieren al(los) microorganismo(s) u otro material biológico depositado mencionado en la descripción en:	
2-1	página	21
2-2	línea	11-24
2-3	Identificación del depósito	
2-3-1	Nombre de la institución de depósito	Colección Española de Cultivos Tipo Universidad de Valencia, Edificio de Investigación, Campus de Burjasot, 46100 Burjasot (Valencia), Spain
2-3-2	Dirección de la institución depositaria	24 Marzo 1999 (24.03.1999)
2-3-3	Fecha de depósito	CECT 5145
2-3-4	Número de orden	
2-4	Indicaciones adicionales	Depósito del Plásmido PROM2-1906-LUC
2-5	Las indicaciones son dadas para los siguientes Estados designados	todos los Estados designados



PCT

2000186

Original (para PRESENTACION) - impreso el 05.07.2000 02:10:05 PM

2-6	Suministro separado de indicaciones Estas indicaciones serán presentadas a la Oficina Internacional posteriormente	NINGUNA
-----	---	---------

## PARA USO DE LA OFICINA RECEPTORA UNICAMENTE

0-4	Este formulario fue recibido con la solicitud internacional: (si o no)	SI'
0-4-1	Funcionario autorizado	Dolores Bautista

## PARA USO DE LA OFICINA INTERNACIONAL UNICAMENTE

0-5	Este formulario fue recibido por la Oficina Internacional el:	
0-5-1	Funcionario autorizado	



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/00/00245

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 : C12Q 1/68, C12N 15.62

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 : C12Q, C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**CIBEPAT, EPODOC, WPI, EMBASE, BIOSIS, MEDLINE, CA, STRAND**

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9837235 A (CORNELL RESEARCH FOUNDATION, INC.) 27 August 1998 (27.08.98) page 4, line 35 – page 5, line 15; page 6, line 9- page 7, line 2; page 8, line 3 – page 9, line 3; example 1; claims 1-9	1-9
Y	the whole document	1-9
X	JAOU-CHEM, H et al. «Gene transfer to cultured human endometrial stromal cells: a model to study cyclooxygenase-2 gene regulation», FERTILITY AND STERILITY, 1998, Vol.70, No. 4, pages 734-739.	1-9
	the whole document	
X	HIROYASU, I. et al. «Transcriptional role of nuclear factor KB site in the induction by lipopolysaccharide and suppression by dexamethasone of cydoxygenase-2 in U937 cell». BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, 1998, Vol. 244, pages 143-148.	1-9
	the whole document	
Y	US 5556754 A (SINGER et al.) 17 September 1996 (17.09.96)	1-9
	the whole document	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

“A” document defining the general state of the art which is not considered to be of particular relevance

“E” earlier document but published on or after the international filing date

“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

“O” document referring to an oral disclosure, use, exhibition or other means

“P” document published prior to the international filing date but later than the priority date claimed

“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

“X” document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

“Y” document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

“&” document member of the same patent family

Date of the actual completion of the international search  
16 October 2000 (16.10.00)

Date of mailing of the international search report  
06 November 2000 (06.11.00)

Name and mailing address of the ISA/  
S.P.T.O.

Authorized officer

Telephone No.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/ES 00/00245

C. (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5445941 A (YANG) 29 August 1995 (17.09.95) the whole document	1-9
A	US 5569588 A (ASHBY et al.) 29 October 1996 (29.10.96) the whole document	1-9

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International Application No  
PCT/ ES 00/00245

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9837235 A	27.08.1998	AU 6015098 A	09.09.1998
US 5556754 A	17.09.1996	US 5871950 A	16.02.1999
		AU 695120 B	06.08.1998
		WO 9428897 A	22.12.1994
		EP 0702554 A	27.03.1996
		CA 2164641 A	22.12.1994
		AU 056494 A	03.01.1995
		JP 8511266 T	26.11.1996
US 5445941 A	29.08.1995	PL 177706 B	31.01.2000
		IL 109990 A	20.06.1999
		NZ 286125 A	24.11.1997
		AU 2871097 A	25.09.1997
		AU 677319 B	17.04.1997
		ZA 9404160 A	13.12.1995
		PL 303915 A	09.01.1995
		HU 70326 A	28.09.1995
		EP 0629697 A	21.12.1994
		CZ 9401475 A	14.06.1995
		CA 2126294 A	22.12.1994
		BR 9402480 A	25.01.1995
		AU 6470194 A	22.12.1994
		NO 942313 A	22.12.1994
		FI 942958 A	22.12.1994
		CN 1102437 A	10.05.1995
		JP 7184661 A	25.07.1995
US 5569588A	29.10.1996	AU 724474 B	21.09.2000
		JP 10507647 T	28.07.1998
		CA 2202154 A	20.02.1997
		EP 0791078 A	27.08.1997
		AU 6720996 A	05.03.1997
		WO 9706277 A	20.02.1997





# INFORME DE BÚSQUEDA INTERNACIONAL

Solicitud internacional n°  
PCT/ ES 00/00245

## A. CLASIFICACIÓN DEL OBJETO DE LA SOLICITUD

CIP<sup>7</sup> C12Q 1/68, C12N 15/62,

De acuerdo con la Clasificación Internacional de Patentes (CIP) o según la clasificación nacional y la CIP.

## B. SECTORES COMPRENDIDOS POR LA BÚSQUEDA

Documentación mínima consultada (sistema de clasificación, seguido de los símbolos de clasificación)

CIP<sup>7</sup> C12Q, C12N

Otra documentación consultada, además de la documentación mínima, en la medida en que tales documentos formen parte de los sectores comprendidos por la búsqueda

Bases de datos electrónicas consultadas durante la búsqueda internacional (nombre de la base de datos y, si es posible, términos de búsqueda utilizados)

CIBEPAT, EPODOC, WPI, EMBASE, BIOSIS, MEDLINE, CA, STRAND

## C. DOCUMENTOS CONSIDERADOS RELEVANTES

Categoría*	Documentos citados, con indicación, si procede, de las partes relevantes	Relevante para las reivindicaciones n°
X	WO 9837235 A (CORNELL RESEARCH FOUNDATION, INC.) 27.08.1998, página 4, línea 35 - página 5, línea 15; página 6, línea 9 - página 7, línea 2; página 8, línea 3 - página 9, línea 3; ejemplo 1; reivindicaciones 1-9	1-9
Y	Todo el documento	1-9
X	JAOU-CHEM, H. et al. "Gene transfer to cultured human endometrial stromal cells: a model to study cyclooxygenase-2 gene regulation", FERTILITY AND STERILITY, 1998, Vol. 70, No. 4. páginas 734- 739 Todo el documento	1-9

☒ En la continuación del recuadro C se relacionan otros documentos ☒ Los documentos de familia de patentes se indican en el anexo

### \* Categorías especiales de documentos citados:

"A" documento que define el estado general de la técnica no considerado como particularmente relevante.

"E" solicitud de patente o patente anterior pero publicada en la fecha de presentación internacional o en fecha posterior.

"L" documento que puede plantear dudas sobre una reivindicación de prioridad o que se cita para determinar la fecha de publicación de otra cita o por una razón especial (como la indicada).

"O" documento que se refiere a una divulgación oral, a una utilización, a una exposición o a cualquier otro medio.

"P" documento publicado antes de la fecha de presentación internacional pero con posterioridad a la fecha de prioridad reivindicada.

"T" documento ulterior publicado con posterioridad a la fecha de presentación internacional o de prioridad que no pertenece al estado de la técnica pertinente pero que se cita por permitir la comprensión del principio o teoría que constituye la base de la invención.

"X" documento particularmente relevante; la invención reivindicada no puede considerarse nueva o que implique una actividad inventiva por referencia al documento aisladamente considerado.

"Y" documento particularmente relevante; la invención reivindicada no puede considerarse que implique una actividad inventiva cuando el documento se asocia a otro u otros documentos de la misma naturaleza, cuya combinación resulta evidente para un experto en la materia.

"&" documento que forma parte de la misma familia de patentes.

Fecha en que se ha concluido efectivamente la búsqueda internacional. 16 Octubre 2000 (16.10.2000)

Fecha de expedición del informe de búsqueda internacional

06 NOV 2000

06. 11. 00

Nombre y dirección postal de la Administración encargada de la búsqueda internacional O.E.P.M.  
C/Panamá 1, 28071 Madrid, España.  
n° de fax +34 91 3495304

Funcionario autorizado

José Luis Vizán

n° de teléfono + 34 1 3495524

# INFORME DE BÚSQUEDA INTERNACIONAL

Solicitud internacional n°

PCT/ ES00/00245

C (Continuación).

## DOCUMENTOS CONSIDERADOS RELEVANTES

Categoría *	Documentos citados, con indicación, si procede, de las partes relevantes	Relevante para las reivindicaciones n°
X	HIROYASU, I. et al. "Transcriptional role of the nuclear factor KB site in the induction by lipopolysaccharide and suppression by dexamethasone of cydoxygenase-2 in U937 cells". BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, 1998, Vol. 244, páginas 143-148. Todo el documento	1-9
Y	US 5556754 A (SINGER et al.) 17.09.1996. Todo el documento	1-9
A	US 5445941 A (YANG) 29.08.1995 Todo el documento	1-9
A	US 5569588 A (ASHBY et al) 29.10.1996 Todo el documento	1-9

**INFORME DE BÚSQUEDA INTERNACIONAL**  
Información relativa a miembros de familias de patentes

Solicitud internacional n°

PCT/ ES 00/00245

Documento de patente citado en el informe de búsqueda	Fecha de publicación	Miembro(s) de la familia de patentes	Fecha de publicación
WO 9837235 A	27.08.1998	AU 6015098 A	09.09.1998
US 5556754 A	17.09.1996	US 5871950 A	16.02.1999
		AU 695120 B	06.08.1998
		WO 9428897 A	22.12.1994
		EP 0702554 A	27.03.1996
		CA 2164641 A	22.12.1994
		AU 056494 A	03.01.1995
		JP 8511266 T	26.11.1996
US 5445941 A	29.08.1995	PL 177706 B	31.01.2000
		IL 109990 A	20.06.1999
		NZ 286125 A	24.11.1997
		AU 2871097 A	25.09.1997
		AU 677319 B	17.04.1997
		ZA 9404160 A	13.12.1995
		PL 303915 A	09.01.1995
		HU 70326 A	28.09.1995
		EP 0629697 A	21.12.1994
		CZ 9401475 A	14.06.1995
		CA 2126294 A	22.12.1994
		BR 9402480 A	25.01.1995
		AU 6470194 A	22.12.1994
		NO 942313 A	22.12.1994
		FI 942958 A	22.12.1994
		CN 1102437 A	10.05.1995
		JP 7184661 A	25.07.1995
US 5569588A	29.10.1996	AU 724474 B	21.09.2000
		JP 10507647 T	28.07.1998
		CA 2202154 A	20.02.1997
		EP 0791078 A	27.08.1997
		AU 6720996 A	05.03.1997
		WO 9706277 A	20.02.1997



# PATENT COOPERATION TREATY

# PCT

REC'D 24 OCT 2001

WIPO PCT

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

16

Applicant's or agent's file reference 2000186	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/ES00/00245	International filing date (day/month/year) 11/07/2000	Priority date (day/month/year) 12/07/1999
International Patent Classification (IPC) or national classification and IPC C12Q1/68		
Applicant LABORATORIOS DEL DR. ESTEVE S.A. et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 7 sheets, including this cover sheet.

- ☒ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 29 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 12/02/2001	Date of completion of this report 19.10.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Herrero, M Telephone No. +49 89 2399 8542 



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/ES00/00245

## I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

### Description, pages:

1-27 as received on 26/09/2001 with letter of 18/09/2001

### Claims, No.:

1-8 as received on 26/09/2001 with letter of 18/09/2001

### Drawings, sheets:

1/10-10/10 filed with the demand

### Sequence listing part of the description, pages:

1-2, filed with the letter of 18.09.01

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:





**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**

International application No. PCT/ES00/00245

- ☐ the description,      pages:  
☐ the claims,      Nos.:  
☐ the drawings,      sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:  
**see separate sheet**

**V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

1. Statement

Novelty (N)	Yes:	Claims	1-8
	No:	Claims	
Inventive step (IS)	Yes:	Claims	1-8
	No:	Claims	
Industrial applicability (IA)	Yes:	Claims	1-8
	No:	Claims	

2. Citations and explanations  
**see separate sheet**

**VII. Certain defects in the international application**

The following defects in the form or contents of the international application have been noted:  
**see separate sheet**

**VIII. Certain observations on the international application**

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:  
**see separate sheet**



## **SECTION I**

### **6. Additional observations**

- 6.1 This preliminary examination report takes into account the content of the Applicants' letter of 18.08.01 in reply to the written opinion dated 09.07.01, as well as the additional technical information enclosed therein.
- 6.2 It has been noticed that in the Applicants' letter dated 18.09.01 a wrong number has been assigned to the present International Patent Application (i.e. PCT/ES 00/00186 instead of the actual PCT/ES 00/00245).

## **SECTION V**

### **2. CITATIONS AND EXPLANATIONS**

- 2.1 The following documents have been considered for the purposes of this report:

- D1: WO 98/37235  
D2: Huang, J-C and M. Y. Dawood (1998) Fertility and Sterility **70**:734-739  
D3: Inoue, H. and T. Tanabe (1998) Biochem. Biophys. Res. Comm. **244**:143-148  
D4: Blanco, J.C.G. et al (19.06.00) J. Exp. Med. **191**:2131-2144

D4 has been cited in the Applicants' letter dated 18.09.01 (a copy of the document is appended hereto).

D4 (published on 19.06.00) is not part of the state of the art according to Rule 64 (1) PCT, as the date of priority of 12.07.99 is hereby validly claimed.

D1 describes a procedure suitable for screening agents as candidates for drugs which suppress induction of COX-2 promoter (see e.g. page 3, lines 7-14; the paragraph bridging pages 4-5 and Examples 1 and 2). This screening procedure



relies on the use a recombinant vector (the plasmid referred to on page 15, line 3 as "COX-2 promoter construct") comprising a construct containing 1432 bases 5' of the human COX-2 transcription start site (-1432/+59) ligated to the luciferase gene. To carry out the subject screening procedure, 1483 squamous carcinoma cells (an apparently conventional cell line in the related art) are transfected with the aforementioned vector containing the construct of interest (see also page 6, lines 3-25 and page 7, lines 27-36 bridging over page 8, lines 1-30 of D1).

D2 describes the determination of the effect of PMA and IL-1 $\beta$  on COX-2 promoter activity (cf paragraph bridging left and right columns on page 736), based on the measurement of luciferase activity generated in endometrial stromal cells transfected with a vector denominated pGL-900. This vector contained a promoter-less luciferase reporter (from pGL-3) and a 900-base pair promoter sequence (-891 to +9 relative to the transcription start site) corresponding to the human COX-2 gene. Among other applications the approach disclosed in D2 can be used to explore the roles of different promoter regulatory elements in COX-2 gene activation (cf page 738, right column, last sentence of the discussion).

D3 studies the transcriptional role of the NF-kB site (nucleotides -223 to -214) of the COX-2 gene in U937 cells employing a luciferase reporter vector driven by the human COX-2 promoter region (nucleotides -327 to +59), mutated at both the cAMP response element and the NF-IL6 site, stably transfected into U937 cells. Attention is drawn to the assays carried out to determine the involvement of the NF-kB site in the suppression of promoter activity by dexamethasone and herbimycin A employing this reporter system described on page 146, left column, second paragraph.

D4 shows that IFN- $\gamma$  works as a pivotal regulator of the Cox-2 gene, activating its expression or coactivating LPS- and IL-1 $\alpha$ -dependent Cox-2 expression in primary murine macrophages. This regulation is dependent on the expression of IRF-1 (interferon regulatory factor-1), and requires the presence of two novel ISREs (interferon stimulated responsive elements) that are localized in the promoter of the murine Cox-2 gene and are conserved in the human Cox-2 gene.



- 2.2 The hereby claimed DNA constructs (Claims 1-2), vector (Claim 3), cell lines (Claims 4-7) and assay method for the search of compounds that selectively inhibit the induction at a transcriptional level of Cox-2 (Claim 8) would appear to relate to novel and inventive subject-matter which satisfies the criteria set forth in Art. 33(2) and (3) PCT for the following reasons:

The DNA construct defined in present Claim 1 comprises the sequence lying between the nucleotides -1796 and +104 of the promoter of the human Cox-2 gene. This DNA construct is therefore novel (Art. 33(2) PCT) over related DNA constructs disclosed in the cited prior art (see above), which respectively comprise the corresponding nucleotide sequences -1445/-1432 to +59 (D1), -891 to +9 (D2) or -327 to +59 (D3).

The desirability of including in a reporter DNA construct of the type hereby claimed nucleotide sequences of the human Cox-2 gene promoter region extending further upstream than position -1475, relative to the transcription start site of the gene, seems to be non-derivable (Art. 33(3) PCT) from the teachings of the prior art (see page 6, lines 3-25 of D1), since the existence of promoter regulatory elements of the Cox-2 gene located further upstream than said position was apparently not known at the date of priority of the present application (see D4).

On the other hand, due to the presence in the hereby characterized reporter construct of an additional regulatory element (located between positions -1796 and -1475) the claimed assay method for the search of compounds which selectively inhibit the induction at a transcriptional level of Cox-2 by a suitable stimulus apparently makes feasible the detection of a wider spectrum of compounds of interest.

The subject-matter encompassed by Claims 1-8 is also susceptible of industrial applicability (Art. 33(4) PCT).





**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/ES00/00245

**SECTION VII**

The words appearing in drawing sheets 1/10, 2/10, 4/10, 5/10 and 6/10 have not been translated.

**SECTION VIII**

The description is not in conformity with the claims as required by Rule 5.1(a)(iii) PCT.



CELL LINE THAT COMPRISES THE PROMOTER OF CYCLOOXYGENASE-2  
(COX-2) AND A REPORTER GENE, AND USE THEREOF IN THE  
SEARCH FOR SELECTIVE INHIBITORS OF THE TRANSCRIPTIONAL  
INDUCTION OF COX-2

5

FIELD OF THE INVENTION

This invention relates, in general, to the search for products with potential therapeutic applications. In particular, the invention relates to a method for the search for compounds that selectively inhibit the induction at a transcriptional level of cyclooxygenase-2 that comprises the use of a cell line that expresses in a stable manner a construct of DNA in which the gene promoter sequence of cyclooxygenase-2 controls the expression of a reporter gene in response to appropriate stimuli.

20

BACKGROUND OF THE INVENTION

The cyclooxygenase (cox) is an enzyme implicated in numerous processes. Two isoforms of cox are known, cyclooxygenase 1 (cox-1) and cyclooxygenase 2 (cox-2). Although both isoforms are related to the production of prostaglandins implicated in physiological processes, it seems that cox-2 is the isoform predominately implicated in various pathologies such as inflammation, carcinogenesis, angiogenesis and certain neurodegenerative processes.

30

Induction at a transcriptional level of cox-2 occurs



in response to several factors, among which can be found the expression of oncogenes, the treatment of tumours with promoters, mytogenous, pro-inflammatory stimuli, growth factors and cytokines [reviewed by Smith and DeWitt, 1996; Griswold and Adams, 1996; Jouzeau et al., 1997 (see section relating to REFERENCES)]. In most cases, the induction of this enzyme translates into an increase in the synthesis of prostaglandins, although other modes of action cannot be discarded.

10

The capacity of certain drugs of the family of non-steroidal anti-inflammatory drugs (NSAIDs) to inhibit cox-2 explains their therapeutic effects [reviewed by Smith and DeWitt, 1996; Griswold and Adams, 1996; Jouzeau et al., 1997]. Similarly, there is growing evidence that the inhibition of cox-2 both by NSAIDs and by glucocorticoids or by cyclosporin A has immunosuppressive effects [Iñiguez et al., 1998; Hall and Wolf, 1997; Zhou et al., 1994; and the reviews cited earlier]. Other actions of induction of cox-2 relate to the implication of this enzyme in cancer, angiogenesis and neurodegenerative processes such as Alzheimer's disease.

It has been found that both inhibition of the induction at a transcription level of cox-2 and the enzymatic inhibition of cox-2 attenuate these processes [Shiff et al., 1996; Tsujii et al., 1997 y 1998; Subbaramiah et al., 1998; Pasinetti, 1998].

After discovering the inducible cox-2 isoform of the cyclooxygenase enzyme, the methods for identification of new anti-inflammatory drugs have focussed on selecting



compounds that are selective inhibitors of the enzymatic activity of cox-2 against the constitutive isoform cox-1.

There are several types of system for this. Some use in vitro assays with purified or semi-purified cox-2  
5 [Famaey, 1997; Noreen et al., 1998]. Other authors use animal or human cell lines that predominantly express the cox-2 isoforms in natural conditions or after induction with stimuli [Famaey, 1997; Berg et al., 1997]. Some use cell lines of animal or human origin in which the cox-2  
10 protein is over-expressed by means of stable transfection of cDNA coding for this protein [Lora et al., 1997; O'Neill et al., 1995; Cromlish and Kennedy, 1996]. In some cases, it has been possible to determine using mRNA analysis whether such compounds inhibit the induction of  
15 cox-2 at a transcriptional level [Tao et al., 1998; Subbaramiah et al., 1998]. Systems have also been established for studying inhibitory compounds by means of in vivo assays, either with whole blood or with purified cells from healthy donors [Famaey, 1997; Brideau et al.,  
20 1996].

In any case, the main limitation of these systems lies in the fact that they allow selection of compounds that inhibit the enzymatic activity of the cox-2 enzyme,  
25 without considering their effects on the induction of the production of the protein, the step prior to production of prostaglandins by this enzyme. In addition to this limitation, it has been shown that the relative potencies of these compounds vary for the same drug for different  
30 types of assay. Similarly, an important aspect for consideration concerns the inhibition of the





physiological activity of cox-2, which would also be inhibited by the type of compounds identified by the aforementioned systems, which could lead to adverse side effects.

5

It is known from WO 98/37235 a method of screening agents as candidates for drugs which suppress induction of COX-2 promoter. This screening procedure relies on the use of a recombinant vector comprising  
10 a construct containg 1432 bases from the COX-2 transcription start site (-1432 to +59) ligated to the luciferase gene.

Document Huang, J-C and M.Y. Dawood (1988)  
15 **Fertility and Sterility** 70:734-739 discloses the determination of the effect of PMA and IL-1 $\beta$  on COX-2 promoter activity based on the measurement of luciferase activity generated in endometirial stromal cells transfected with a vector that contains a  
20 luciferase reporter and a 900-base pair promoter sequence (-891 to +9 relative to the transcription site) corresponding to the human gene.

Document Inoue, H. and T. Tanabe (1998) **Biochem.**  
25 **Biophys. Res. Comm.** 244: 143-148 studies the transcriptional role of the NF-kB site of the COX-2 gene in U937 cells employing luciferase reporter vector driven by the human COX-2 promoter region (nucleotides -327 to +59) stably transfected into U937  
30 cells.



There is therefore a need to develop a method for searching for compounds that selectively inhibit induction of cox-2 at a transcriptional level by different stimuli that overcomes the drawbacks mentioned  
5 above.

#### OVERVIEW OF THE INVENTION

This invention provides a solution to the existing need that consists of developing an assay system for the  
10 search of compounds that selectively inhibit the induction of cox-2 at a transcriptional level by different stimuli. This criterion allows compounds to be selected that inhibit the production of cox-2, and so will act as inhibitors of the actions derived from an  
15 increase in the expression of cox-2 and of the subsequent increase in the production of prostaglandins that set off various pathological processes. Among other processes, the following processes can be highlighted: inflammatory processes, uncontrolled cellular proliferation,  
20 angiogenesis, carcinogenesis and neurodegenerative pathologies, as were described earlier. The criterion for the selection of compounds according to this invention lies in the inhibition of the inducible activity of the promoter of cox-2. Therefore, those  
25 compounds that inhibit the physiological basal activity of production of cox-2 will not be selected.

For the development of the solution provided by this invention, it has been necessary to construct a cell line  
30 that expresses in a stable fashion a construct of DNA in which the promoter sequence of the cox-2 gene controls



the expression of the reporter gene in response to a suitable stimulus. The regulation of the expression of the cox-2 gene is determined by the regulatory activity of its promoter, while the measurement of the activity of the reporter gene provides an indirect measure of the activity of the promoter of cox-2 in response to different agents.

Thus, an object of this invention constitutes a DNA construct (or recombinant DNA) that comprises a promoting sequence of the cox-2 gene and a reporter gene, operatively joined to each other, such that said promoter sequence of the gene controls the expression of the reporter gene in response to a suitable stimulus.

15

An additional object of this invention constitutes a vector, such as a plasmid or an expression vector, that comprises said DNA construct.

Another additional object of this invention constitutes a cell line that contains said DNA construct, or said plasmid that contains said DNA construct, that expresses it in a stable fashion.

Finally, another additional object of this invention constitutes a method for the search for compounds that selectively inhibit the induction of cyclooxygenase-2 at a transcriptional level, that comprises the use of said cell line that expresses said DNA construct in a stable fashion.



## BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the sequence of the promoter zone of the *cox-2* gene. The arrows indicate the sequences of hybridisation of the oligonucleotides used in the PCR reaction [polymerase chain reaction].

Figure 2 shows the strategy for cloning the promoting region of the *cox-2* gene in the pXP2 plasmid in order to obtain the construct prom2-1906-LUC.

Figure 3 shows the results of the analysis by RT-PCR [reverse transcription-polymerase chain reaction] of the expression of mRNA of *cox-2* in Jurkat cells. In Figure 3(A) the effect of treatment with PMA and PMA + calcium ionophore A23187, hereinafter PMA-Ion, on the expression of mRNA of *cox-2* is shown. In Figure 3(B) the inhibition by cyclosporin of the transcriptional induction of *cox-2* is shown. In both cases, the result obtained for the non-inducible mRNAs of the *cox-1* isoform and the glycerol-aldehyde dehydrogenase (GAPDH) by way of control is shown.

Figure 4 shows the result of the stimulation by PMA or by PMA+Ion of the luciferase activity in Jurkat cells transitorily transfected with the prom2-1906-LUC construct. As a control, it is checked that both the prom1-898-LUC construct and the empty plasmid PXP2 are not inducible.

Figure 5 shows the result of inhibition by





cyclosporin A (CsA) of the stimulation caused by PMA-Ion of the construct prom2-1906-LUC in the transient transfection in Jurkat cells.

5        Figure 6 shows the results of an experiment of transitory transfection with the prom2-1906-LUC construct and treatment with dexamethasone.

10        Figure 7 shows the results of the luciferase activity of different clones obtained after stable transfection with the prom2-1906-LUC construct.

15        Figure 8 shows the results of inhibition by cyclosporin A (CsA) of the stimulation by PMA+Ion of the luciferase activity of the stable clones of Jurkat-1906LUC.

20        Figure 9 shows the results obtained for the inhibition by the glucocorticoide dexamethasone (Dex) on the stimulation by PMA+Ion of the luciferase activity of the stable clones of Jurkat-1906LUC.

25        Figure 10 shows the inhibition by Resveratrol (Res) of the induction of luciferase activity of the stable clones as a control of the assay system.

#### DETAILED DESCRIPTION OF THE INVENTION

30        This invention provides a DNA construct (or recombinant DNA) that comprises the whole part of a promoter sequence of the cyclooxygenase 2 (cox-2) gene



and a reporter gene, operatively joined to each other, such that said promoter sequence of the cox-2 gene controls the expression of said reporter gene in response to a suitable stimulus.

5

The promoter sequence of the cox-2 gene may be of any origin, although preferably said sequence will proceed from the human cox-2 gene.

10

As a reporter gene, any reporter gene may be used that is able to produce an easily detectable signal, chosen from among those normally used in these types of transfection assays, for example, the chloranphenicol acetyl transferase (CAT) gene, the beta galactosidase ( $\beta$ -gal) gene, the luciferase gene, for example, from glow-worm or from *Renilla*. In a particular embodiment of this invention, said reporter gene is the luciferase gene of glow-worm because of its extreme sensitivity, speed and ease of use and low cost of the assay for its detection.

15  
20

The invention also provides a vector, such as a plasmid or an expression vector, which contains the aforementioned DNA construct. In principle, any vector can be used that is suitable for inserting into said DNA construct. These vectors are useful for the transformation of cells.

The cell line provided by this invention comprises, and expresses in a stable fashion, said DNA construct that comprises all or part of a promoter sequence of the cox-2 gene and a reporter gene, operatively joined to

25  
30



each other, such that said promoter sequence of the cox-2 gene controls the expression of said reporter gene in response to a suitable stimulus.

5           The transformed cell line that contains the  
aforementioned DNA construct may originate from any  
suitable cell line able to express said DNA construct in  
a stable fashion, for example, a cell line of human  
origin such as a line of cells of the T-lymphocyte type,  
10   Hep-G2 cells derived from hepatocellular carcinoma, Hela  
cells derived from an adenocarcinoma of the cervix, cells  
of monocyte-macrophage type, for example, the lines U937  
and THP-1, etc. In a particular embodiment of this  
invention, a Jurkat cell line has been selected  
15   (originally described by Schneider et al., 1977) as a  
representative example of a transformed cell line of the  
T-lymphocyte type as a model for studying the expression  
of a gene related to the immune response. In addition,  
said cell line is easy to grow and provides a high yield  
20   of cells per unit time and volume (ml) of culture.

The cell line provided by this invention can be used  
as an assay system in the search and identification  
(screening) of compounds that selectively inhibit the  
25   induction at a transcriptional level of cox-2 by  
different stimuli.

The cell line provided by this invention can be  
easily obtained using conventional procedures of Genetic  
30   Engineering, for example, by means of a process that  
comprises (i) the isolation of a promoter sequence of the



cox-2 gene, (ii) the cloning of said sequence in a vector that contains the reporter gene, in a position in which said promoter sequence is able to control the expression of said reporter gene, and (iii) the transfection of a  
5 suitable cell line with said plasmid. In Example 3, there is described in detail a specific way for obtaining individual clones of transformed Jurkat cells which express the reporter gene (luciferase) in a stable fashion, denominated Jurkat-C3-1906LUC, Jurkat-F9-1906LUC  
10 and Jurkat-C7-1906LUC, in which the basal luciferase activity was determined and it was checked that the expression of the reporter gene (luciferase) was being induced in response to the same stimuli as the promoter of cox-2 as had been previously established with  
15 transitorily transfected cells.

The assay system (cell line) provided by this invention has been previously validated by the transitory transfection of the prom2-1906-LUC construct and the  
20 analysis of the luciferase activity under different stimuli and inhibitors. The results obtained were compared with a non-inducible control of a similar construct in which, instead of the promoter of cox-2, the promoter of the cox-1 isoform was put in place and with  
25 the empty vector pXP2. Under the same conditions, the behaviour of the endogenous cox-2 gene was also determined by experiments of RT-PCR in which the expression of mRNA is analysed. As a non-inducible control, the behaviour of the endogenous gene of the cox-  
30 1 isoform and the non-inducible gene of glycerol-aldehyde-phosphate dehydrogenase (GAPDH) was determined.





The main treatments were activators such as phorbol ester PMA (10 ng/ml) and the combination of PMA and calcium ionophore A23187 [PMA+Ion] (1  $\mu$ M). Treatment with drugs that inhibit the induction by PMA+Ion were carried out  
5 with dexamethasone (1  $\mu$ M), and with cyclosporin A (100 ng/ml).

Example 2 includes some validation assays of the assay system provided by this invention in transitory  
10 transfection, as well as the expression of the endogenous genes in Jurkat cells. Also, in Examples 3 and 4, some assays are presented carried out with the stable clones obtained, with compounds that induce and inhibit the induction the cox-2 promoter.

15

The assay system provided by this invention is useful for the search and identification (screening) of compounds that selectively inhibit the induction at a transcriptional level of cox-2 by different stimuli.  
20 This type of selective inhibitor of cox-2 may have numerous potential therapeutic applications as the implications derived from the induction of cox-2 not only affect the inflammatory response, but also processes related to uncontrolled cellular proliferation and  
25 formation of tumours (for example, the appearance of adenomas, cancer of the colon and the development of polyps and angiogenesis, among others), with immunosuppressant actions and with neurodegenerative processes such as Alzheimer's disease. As a result, it  
30 might be supposed that the compounds that selectively inhibit the transcriptional induction of cox-2 may be



useful as anti-inflammatory agents, as compounds able to attenuate uncontrolled cellular proliferation and/or the process of tumorigenesis, as immunosuppressants or as potential drugs with therapeutic properties in  
5 Alzheimer's disease.

The invention also provides an assay method for searching and identifying (screening) compounds that selectively inhibit the induction at a transcriptional  
10 level of cox-2 by a suitable stimulus (described in Example 4) which comprises bringing the cell line provided by this invention (assay system) into contact with the compound to be assayed, in other words, with the compound whose potential selective inhibitory activity of  
15 the induction at a transcriptional level of cox-2 it is wanted to test, in conditions that allow the transcription of cox-2, and detecting, and/or measuring, the signal indicative of the expression of activity due to the reporter gene. Alternatively, if so desired, the  
20 assay method object of this invention can be performed by bringing the cell line, the assay system and the compound that activates transcription induction of cox-2 into contact.

25 In the assay method provided by this invention, the regulation of the expression of the cox-2 gene is determined by the regulatory activity of its promoter, while the activity of the reporter gene provides an indirect measure of the activity of the cox-2 promoter in  
30 response to different agents.



The assay method for the search and identification of compounds that selectively inhibit the induction at a transcriptional level of cox-2 by an appropriate stimulus provided by this invention allows the selection of compounds that inhibit the production of cox-2 by means of a criterion based on the inhibition of the inducible activity of the cox-2 promoter, which will not select those compounds that inhibit the basal physiological activity of the production of cox-2.

10

The following examples serve to illustrate preferred embodiments of the invention, but they should not be considered as limiting the scope thereof.

15

#### EXAMPLE 1

**Production of a DNA construct that comprises a promoter sequence of cox-2 and the luciferase gene**

##### 1.1 Cloning the promoter of cox-2

In the first place, the promoter sequence of the human cox-2 gene was cloned from the sequence described by [Tazawa et al., 1994] represented in Figure 1.

The technique of polymerase chain reaction (PCR) was used, with the initiating oligonucleotides or "primers" designed for selective amplification of the fragment of DNA corresponding to the promoter sequence of this gene.

The template DNA used was genomic DNA from the Jurkat human lymphocyte cell line. The oligonucleotides used were those identified as SEC.ID.No.: 1 and SEC.ID.No.: 2 [see the section concerning the LIST OF SEQUENCES].

30



These oligonucleotides amplify a sequence that ranges from nucleotide -1796 to nucleotide +104 of the promoter zone of the cox-2 gene (see Figure 1). For the PCR reaction, the Advantage cDNA PCR kit [Clontech] was used with 30 cycles repeated every 45 seconds at 94°C and 3 minutes at 68° C in a PTC-200 thermocycler PTC-200 [MJ Research].

#### 10           1.2 Construction of the expression vector

The fragments generated after amplification were subcloned into the plasmid pXP2 [Nordeen, 1988] which contains the sequence that codes for the luciferase gene which will be used as the reporter gene (see Figure 2).

15   The oligonucleotides were designed such that at the 5' end they contain an additional recognition sequence for restriction enzymes. After amplification, double chain ends are generated that contain the restriction targets BamHI at the 5' end and BglIII at the 3' end. The pXP2

20   vector contains a BglIII target at the multiple cloning site, which generates ends compatible both with BglIII and BamHI ends. After digestion of the insert obtained by PCR containing the promoting sequence with the BglIII and BamHI enzymes and the pXp2 vector with BglIII, the binding

25   of these sequences was performed. In this fashion, the plasmid prom2-1906-LUC was obtained in which the sequence (-)1796-(+)104 of the cox-2 promoter is located in front of the luciferase gene, controlling its expression. This construction was sequenced to check the fidelity of the

30   promoter sequence and to verify the cloning site.





## Example 2

Experiments in analysis of the regulation of the activity of the promoter of cox-2 by means of experiments with RT-PCR and transitory transfection of the prom2-1906-LUC construct in the Jurkat cell line

In order to study the regulation of the expression of the endogenous cox-2 gene in the Jurkat cell line, the analysis of the expression of its mRNA was performed by experiments with RT-PCR. At the same time, in order to validate the prom2-1906-LUC construct and to check that the regulation of the cloned promoter is as expected, experiments were carried out of transitory transfection of the prom2-1906-LUC construct using the Jurkat cell line. In both experiments, the cells were treated with compounds that stimulate and inhibit the induction of the promoter of cox-2.

Treatment with activator compounds were carried out using the phorbol ester PMA (Phorbol 12-Myristate 13-Acetate) (10 ng/ml) (Sigma) and the combination of PMA and Calcium ionophore A23187 (1 $\mu$ M) (Sigma), hereinafter PMA+Ion.

Treatment with drugs that inhibit the induction by PMA+Ion were carried out with cyclosporin A (CsA) (100 ng/ml) or the synthetic glucocorticoids Dexamethasone (Dex) (1 $\mu$ M) (Sigma).

2.1 Regulation of the expression of cox-2 mRNA  
in Jurkat cells



The results obtained from the analysis by RT-PCR of the expression of mRNA of cox-2 in Jurkat cells are presented in Figure 3, where the following can be observed:

- 5           a) treatment with PMA (10 ng/ml) produces a slight increase in the expression of the mRNA of cox-2, while the combination treatment with PMA+Ion leads to a greater increase in the expression of this gene at a transcriptional level [Figure 3(A)]; and
- 10           b) inhibition by CsA (100 ng/ml) of the transcriptional induction of cox-2 [Figure 3(B)].

As a control in both cases, the result obtained for the non-inducible mRNAs of the cox-1 isoform and for the  
15   glycerol aldehyde phosphate dehydrogenase (GAPDH) are shown.

## 2.2 Evaluation of the luciferase activity under different stimuli

20           The luciferase activity in Jurkat cells transitorily transfected with the prom2-1906-LUC construction using the Lipofectin agent [Life Technologies] has been analysed. The cells were stimulated with PMA (10 ng/m) (Sigma) and with the combination PMA+Ion (1  $\mu$ M) (Sigma).

25

As a non-inducible control, a similar construct was used in which, instead of the cox-2 promoter, the promoter of the cox-1 isoform was put in place [prom1-898-LUC] and with the empty vector pXP2.

30

The luciferase activity was determined using the kit



"Luciferase Assay System" [Promega], with  $1 \times 10^6$  cells which were lysed in 50  $\mu$ l of lysis buffer. In the extracts obtained, the light emission produced was measured using a luminometer Monolight 2010 [Analytical  
5 Luminiscence Laboratory] with an automatic injection system of 100  $\mu$ l of reagent.

The results obtained are shown in Figure 4. As can be seen in said figure, the results obtained in this  
10 assay are comparable to those obtained in the analysis of mRNA of cox-2 [Example 2.1], in other words, treatment with PMA+Ion produces a greater increase in the number of times there is induction of the luciferase activity (approximately 12 times the basal value). These data  
15 show that the cloned promoter sequence behaves in a similar fashion to the endogenous gene. As a control, it is checked that neither the prom1-898-LUC construct nor the empty plasmid pXP2 are inducible.

20 2.3 Evaluation of the luciferase activity under different inhibitors

The inhibition of stimulation by PMA (10 ng/ml) (Sigma) or PMA+Ion (1  $\mu$ M) (Sigma) on the luciferase activity in Jurkat cells transitorily transfected with  
25 the prom2-1906-LUC construct has been investigated by using the immunosuppressant drug cyclosporin A (CsA) (100 ng/ml). The results obtained are shown in Figure 5 where it can be seen that treatment with CsA (100 ng/ml) reduces stimulation of the promoter of cox-2 in response  
30 to PMA+Ion to values similar to basal ones.



Similarly, the inhibition of stimulation by PMA+Ion (1  $\mu$ M) (Sigma) on the luciferase activity has been analysed in Jurkat cells transitorily transfected with the prom2-1906 construct by means of the glucocorticoide dexamethasone (1  $\mu$ M) (Sigma). The results obtained are shown in Figure 6 where it can be seen that treatment with dexamethasone (1  $\mu$ M) (Dex) reduces stimulation of the promoter of cox-2 in response to PMA+Ion.

10

### Example 3

Production of a cell line that stably expresses a DNA construct that comprises a promoter sequence of cox-2 and the luciferase gene.

For the creation of a cell line that stably expresses the prom2-1906-LUC construct, Jurkat cells were co-transfected with the vector prom2-1906-LUC and a vector denominated pcDNA3.1/Hygro (Invitrogen) which contains the gene for resistance to hygromycin. The transfection was carried out by means of the technique of electroporation in cuvettes of 0.4 cm (BioRad) with  $15 \times 10^6$  cells in 0.5 ml of complete medium [RMPI medium supplemented with 10% of foetal serum, L-glutamine 2mM and a mixture of antibiotics] (All these products were acquired from Life Technologies). The cells were incubated over ice for 10 minutes with 25  $\mu$ g of plasmid prom2-1906-LUC and 5  $\mu$ g of the vector pcDNA3.1./Hygro. After this period, the cells were electroporated in a Gene Pulser II (BioRad) apparatus at 1.500  $\mu$ Faradays of capacitance and a current of 280 V. Next, the cells were incubated over ice for 10 minutes before adding 10 ml of





complete medium. The cells were cultured in this medium in 75 cm<sup>2</sup> culture flasks (Nunc) for 48 hours in a cell incubator at 37° C with a humidity of 95% and 5% of CO<sub>2</sub>. At this time, the medium was changed for complete medium with no antibiotics to which hygromycin (Boehringer Mannheim) was added at a concentration of 200 µg/ml. The cells were cultured in this medium for 30 days with successive changes of medium. During this period the resistant population that survived treatment with the selective antibiotic was selected, in other words, the cells stably transfected with the gene for resistance to the hygromycin antibiotic. In this population the expression of the luciferase gene was analysed in order to determine the presence of transfectants stable for the plasmid prom2-1906-LUC. For this, 1x10<sup>6</sup> cells were lysed in 50 µl of lysis buffer (Promega) and with the extracts obtained, the luciferase activity was determined using the reagents contained in the kit of the "Luciferase Assay System" [Promega]. Measurement of the light produced was determined using a luminometer MonoLight 2010 (Analytical Luminiscence Laboratory) with a system of automatic injection of 100 µl of reagent.

From this polyclonal population (Jurkat-pool-1906LUC) a limit dilution was performed on 96-well plates in complete medium with hygromycin in order to obtain individual clones that expressed the luciferase gene in a stable fashion. These clones were grown until obtaining at least 1x10<sup>6</sup> cells with which measurement of the luciferase activity could be performed as described earlier. In this way, three individual clones were



obtained denominated Jurkat-C3-1906LUC, Jurkat-F9-1906LUC and Jurkat-C7-1906LUC. In these clones the basal luciferase activity was determined in RLU (relative luminescence units) and it was checked that the expression of the luciferase reporter was being induced in response to the same stimuli as the promoter of cox-2 as had been established previously with cells transitorily transfected (see Figure 7). In the three clones, the basal values of luciferase activity increase from 3 to 6 times with a treatment of 6 hours with the phorbol ester PMA and up to 10 - 20 times with a combined treatment PMA+Ion, in a similar fashion to the results obtained in the cells transfected transitorily.

15

#### Example 4

Establishment of an assay system for compounds that regulate the expression of the gene cox-2 in the clones of the cell line that stably expresses a DNA construct that comprises a promoter sequence of cox-2 and the luciferase gene.

20

The clones were cultured on plates of 96-wells at a density of  $1 \times 10^5$  cells in 200  $\mu$ l of RPMI medium supplemented with 2% of foetal serum, L-Glutamine 2 mM and a mixture of antibiotics. The cells were treated for 6 hours with different concentrations of compounds whose activity it was hoped to analyse. In the case of the assay of the activity of these compounds on the induction of activity of the cox-2 promoter, the cells were treated with PMA+Ion for 5 hours, after 1 hour of pre-treatment with the compound to test. After this period, the cells

25  
30



were lysed in 50 µl of lysis buffer and their luciferase activity determined using 20 µl in a luminometer as was described in Example 2.2. Next, some results obtained are shown with compounds previously described as inhibitors  
5 of the stimulation of the cox-2 promoter.

Figure 8 shows the results obtained with the compound cyclosporin A (CsA) which produces an inhibition of the stimulation obtained with PMA+Ion in stable Jurkat  
10 clones, in a similar fashion to that already described (Iñiguez et al., 1998), and that observed in the transitory transfections illustrated in Example 2.3.

Figure 9 shows the results obtained with the  
15 compound Dexamethasone (Dex), which, like glucocorticoids and anti-inflammatories, produces an inhibition of the stimulation obtained with PMA+Ion in the stable Jurkat clones, corresponding to that already described (Smith and DeWitt, 1996) and that observed previously in the  
20 transitory transfections of Example 2.3.

Figure 10 shows the results obtained with the compound Resveratrol (Res), recently described as an inhibitor of the stimulation by PMA of the gene cox-2  
25 (Subbaramiah, et al., 1998). This compound produces an inhibition of the stimulation obtained with PMA+Ion on the stable Jurkat clones.

The validity of the assay system is thus  
30 demonstrated due to its similar behaviour for the clones obtained, for the transitory transfections and the



results obtained with endogenous mRNA. With the use of compounds whose activity on the cox-2 promoter is known, whether they be stimulatory or inhibitory, it is established that it is possible to detect both positively and negatively the basal or induced expression of the cox-2 gene with the assay system developed in present invention.

#### DEPOSIT OF BIOLOGICAL MATERIAL

10

A sample of a Jurkat cell line, denominated J-1906-F9, that stably expresses a DNA construct that comprises a promoter sequence of the cox-2 gene and the luciferase gene, has been deposited in the European Collection of Animal Cell Cultures (ECACC) [Salisbury, United Kingdom] on the 24 March 1999 and has been assigned the access number ECACC 99032405.

A sample of the plasmid prom2-1906-LUC, inserted into *Escherichia coli* DH5, denominated DH5 prom2-1906-LUC, has been deposited in the Spanish Collection of Culture Types (CECT) [Burjassot, Valencia] on the 24 March 1999 and has been assigned the access number CECT 5145.

25

#### REFERENCES

Berg, J., Christoph, T., Widerna, M., and Bodenteich, A. 1997. Isoenzyme-specific cyclooxygenase inhibitors: a whole cell assay system using the human erythroleukemic cell line HEL and the human monocytic





cell line Mono Mac 6. *J. Pharmacol. Toxic Methods.* 37:  
179-86.

Brideau, C., Kargman, S., Liu S., Dallob A.L.,  
5 Enrich E.W., Rodger, I.W., And Chan C.C. 1996. A human  
whole cell assay for clinical evaluation of biochemical  
efficacy of cyclooxygenase inhibitors. *Inflamm. Res.* 45:  
68-74.

10 Cromlish W.A., and Kennedy, B.P. 1996. Selective  
inhibition of cyclooxygenase-1 and -2 using intact insect  
cell assays. *Biochem. Pharmacol.* 52: 1777-85.

Famaey, J.P. 1997. In vitro and in vivo  
15 pharmacological evidence of selective cyclooxygenase-2  
inhibition by nimesulide: an overview. *Inflamm. Res.* 46:  
437-446.

Griswold, D.E., and Adams, J.L. 1996. Constitutive  
20 cyclooxygenase (cox-1) and inducible cyclooxygenase (cox-  
2): Rationale for selective inhibition and progress to  
date. *Med. Res. Rev.* 16: 181.

Hall, V.C., and Wolf, R.E. 1997. Effect of Tenidap  
25 and nonsteroidal antiinflammatory drugs on the response  
of cultured human T cells to interleukin 2 in rheumatoid  
arthritis. *J. Rheumatol.* 24:1467.

Iñiguez, M.A., Punzón, C., and Fresno, M. 1998.  
30 Induction of cyclooxygenase-2 on activated T lymphocytes;  
regulation of T cell activation by cyclooxygenase-2



inhibitors. Sent to publish.

Jouzeau, J.Y., Terlain, B., Abid, A., Nedelec, E.,  
and Netter, P. 1997. Cyclooxygenase isoenzymes. How  
5 recent findings affect thinking about nonsteroidal anti-  
inflammatory drugs. *Drugs* 53: 563.

Lora, M., Morisset, S., Menard, H.A., Leduc, R., and  
de Brum-Fernandes, A.J. 1997. Expression of recombinant  
10 human cyclooxygenase isoenzymes in transfected COS-7  
cells in vitro and inhibition by tenoxicam, indomethacin  
and aspirin. *Prostaglandins Leukot. Essent. Fatty Acids*.  
56: 361-7.

15 Nordeen, S.K. 1988. Luciferase reporter gene vectors  
for analysis of promoters and enhancers. *Biotechniques* 6:  
454-457.

Noreen, Y., Ringbom, T., Perera, P., Danielson, H.,  
20 And Bohlin, L. 1998. Development of a radiochemical  
cyclooxygenase-1 and -2 in vitro assay for identification  
of natural products as inhibitors of prostaglandin  
biosynthesis. *J. Nat. Prod.* 61: 2-7.

25 O'Neill, G.P., Kennedy, B.P. Mancini, J.A., Kargman,  
S., Ouellet, M., Yergey, J., Falgueyret, J.P., Cromlish,  
W.A., Payett, P., Chan, C.C. et al. 1995. Selective  
inhibitors of Cox-2. *Agents Actions Suppl.* 46: 159-68.

30 Pasinetti, G.M. 1998. Cyclooxygenase and  
inflammation in Alzheimer's disease. *Experimental*



approaches and clinical interventions. *J. Neurosci. Res.* 54: 1-6.

Schneider U., Schwenk, H.U., and Bornkamm, G. 1977.  
5 Characterization of EBV-genome negative "null" and "T"  
cell lines derived from children with acute lymphoblastic  
leukemia and leukemic transformed non-Hodkin lymphoma.  
*Int. J. Cancer*, 19:521-6.

10 Sheng, H., Shao, J., Kirkland, S.C., Isakson, P.,  
Coffey, R.J., Morrow, J., Beauchamp, R.D., and DuBois,  
R.N. 1997. Inhibition of human colon cancer cell growth  
by selective inhibition of cyclooxygenase-2. *J. Clin.*  
*Invest.* 99: 2254-2259.

15 Shiff, S.J., Koutsos, M.I., Qiao, L., and Rigas, B.  
1996. Nonsteroidal antiinflammatory drugs inhibit the  
proliferation of colon adenocarcinoma cells: effect on  
cell cycle and apoptosis. *Exp. Cell Res.* 222, 179-188.

20 Smith, W.L., and DeWitt, D.L. 1996. Prostaglandin  
Endoperoxide H Synthase-1 and -2. *Adv. Immunol.* 62:167.

Subbaramiah, K., Chung, W.J., Michualart, P.,  
25 Telang, N., Tanabe, T., Inoue, H., Jang, M., Pezzuto,  
J.M., and Dannenberg, A.J. 1998. Resveratrol inhibits  
cyclooxygenase-2 transcription and activity in phorbol-  
ester treated human mammary epithelial cells. *J. Biol.*  
*Chem.* 273: 21875-882.

30 Tao, X., Schulze-Koops, H., Ma, L., Cai, J., Mao,



Y., and Lipsky, P.E. 1998. Effects of *Tripterygium wilfordii* hook extracts on induction of cyclooxygenase 2 activity and prostaglandin E2 production. *Arthritis Rheum.* 41: 130-8.

5

Tazawa, R., Xu, X.M., Wu, K.K., and Wang, L.H. 1994. Characterization of the genomic structure, chromosomal location and promoter of human prostaglandin H synthase-2 gene. *Biochem. Biophys. Res. Comm.* 203:190-194.

10

Tsujii M., Kawano, S., Tsuji, S., Sawaoka, H., Matsatsugu, H., and DuBois, R.N. 1998. Cyclooxygenase regulates angiogenesis induced by colon cancer cells. *Cell* 93: 705-716.

15

Tsujii, M., Kawano, S., and DuBois, R.N. 1997. Cyclooxygenase-2 expression in human colon cancer cells increases metastatic potential. *Proc. Natl. Acad. Sci. USA* 94: 3336-40.

20

Zhou, L., Ritchie, D., Wang, E.Y., Barbone, A.G., Argentier D., And Lau, C.Y. 1994. Tepoxalin, a novel immunosuppressive agent with a different mechanism of action from cyclosporin A. *J. Immunol.* 153:5026.





## CLAIMS

1. A DNA construct that comprises the sequence lying between the nucleotide (-) 1796 and the nucleotide (+)104  
5 of the promoter of human cyclooxygenase 2 (cox-2) gene and a reporter gene, operatively joined to each other, such that said promoter sequence of the cox-2 gene controls the expression of said reporter gene in response to a suitable stimulus.

10

2. A construct according to claim 1, in which said reporter gene is selected from the luciferase gene, the chloramphenicol acetyltransferase gene and the gene of beta galactosidase.

15

3. A vector that comprises a DNA construct according to any of claims 1 to 2.

4. A cell line that comprises a construct according  
20 to any of claims 1 to 2 or transformed with a vector according to claim 3.

5. A cell line according to claim 4, in which said cell line is derived from a cell line of human origin.

25

6. A cell line according to claim 5, in which said cell line of human origin is a line of Jurkat cells.

7. A cell line according to claims 4 to 6 which  
30 expresses in stable fashion the DNA construct of claims 1 to 2.



8. An assay method for the search for compounds that selectively inhibit the induction at a transcriptional level of cyclooxygenase-2 by a suitable stimulus, that  
5 comprises bringing a cell line according to any of claims 5 to 7, into contact with a compound whose potential selective inhibitory activity of induction at a transcriptional level of cox-2 it is wanted to assay, in conditions that allow the transcription of cox-2, and  
10 detecting, and/or measuring, the signal indicative of the expression of activity due to the reporter gene.



# PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner  
US Department of Commerce  
United States Patent and Trademark  
Office, PCT  
2011 South Clark Place Room  
CP2/5C24  
Arlington, VA 22202  
ETATS-UNIS D'AMERIQUE  
in its capacity as elected Office

<b>Date of mailing</b> (day/month/year) 20 April 2001 (20.04.01)	
<b>International application No.</b> PCT/ES00/00245	<b>Applicant's or agent's file reference</b> 2000186
<b>International filing date</b> (day/month/year) 11 July 2000 (11.07.00)	<b>Priority date</b> (day/month/year) 12 July 1999 (12.07.99)
<b>Applicant</b> FRESNO ESCUDERO, Manuel et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:  
12 February 2001 (12.02.01)

☐ in a notice effecting later election filed with the International Bureau on:  
\_\_\_\_\_

2. The election ☒ was

☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<p>The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland</p> <p>Facsimile No.: (41-22) 740.14.35</p>	<p>Authorized officer</p> <p>Pascal Piriou</p> <p>Telephone No.: (41-22) 338.83.38</p>
--	--

